

Fourier integral operators with open umbrellas and seismic inversion for cusp caustics

Raluca Felea and Allan Greenleaf

Abstract

In general the composition of Fourier integral operators (FIOs) need not be an FIO. Motivated by the problem of linearized seismic inversion in the presence of cusp caustics for the background sound speed, we consider FIOs whose canonical relations have certain two-sided cusp degeneracies, and show that the resulting compositions have wave-front relations in the union of the diagonal and an open umbrella, the simplest type of singular Lagrangian manifold.

1 Introduction

A fundamental problem concerning Fourier integral operators (FIOs) is that, outside of the standard transverse [19] and clean intersection [5, 32] calculus, a composition of two FIOs is typically not another FIO. Describing the operators resulting from the composition of completely general FIOs and placing them in a usable class, with a symbol calculus, Sobolev space estimates, and the possibility of constructing parametrices under suitable ellipticity conditions, is at this point a distant prospect. Some progress has been made for specific geometries arising in integral geometry [16] and inverse problems [6, 7, 21, 28]. In all of these works, the compositions were shown to belong to an existing class, the *pseudodifferential operators with singular symbols*, which are not FIOS and were introduced to construct parametrices for operators of real or complex principal type [24, 18, 26]. Such operators have wave front relation, i.e., the wave front set of the Schwartz kernel, contained in the union of the diagonal and another smooth canonical relation which intersects the diagonal cleanly.

In the present paper, motivated by a linearized inverse problem from seismology for the acoustic wave equation, we analyze FIOs having a special type of degeneracy, which we call a *flat two-sided cusp*, and show that their composition results in operators which are fundamentally more singular than those in works referred to above.

FIOs with *two-sided cusps*, i.e., those for which the projections from the canonical relation both to the left and the right are cusps, arise naturally when considering generalized Radon transforms. They have previously been studied in terms of their L^2 Sobolev regularity properties; it was shown in [3, 13] that there is a loss of $1/4$ derivative compared with the nondegenerate case of local canonical graphs. $L^2 \rightarrow L^q$ estimates for FIOs with one-sided cusps have also been obtained, with T^*T making an appearance through Strichartz estimate type arguments [12]; however, such compositions have not been studied in their own right. We show that for the subclass of *flat two-sided cusps* which we formulate, the normal operator F^*F is not a standard pseudodifferential operator with singular symbol, as it is in the case of fold caustics [28, 6, 7]. This includes the canonical relations underlying the linearized forward scattering operators F in the seismology problem in the presence of caustics of *cusp* type. While there is still a large part of its wave front relation, $WF(K_{F^*F})'$, which is contained in the diagonal, Δ , the remaining portion, $\tilde{C}$, is now no longer a *smooth* canonical relation. Rather, $\tilde{C}$ has the structure of the simplest kind of singular Lagrangian manifold, an *open umbrella* [9]. The diagonal and this open umbrella intersect in codimension one, and we quantize the variety $\Delta \cup \tilde{C}$ by associating to it a class of generalized Fourier integral operators, $I^m(\Delta \cup \tilde{C})$. This is in the spirit of the paired Lagrangian (or $I^{p,l}$) distributions associated to a pair of cleaning intersecting smooth Lagrangians; however, rather than nondegenerate phase functions and product type symbols as in [24, 18, 26], we use a combination of a degenerate phase function (whose gradient exhibits normal crossings) and standard symbols. We remark that operators with wave front relation in the union of Δ and a canonical relation having a conical singularity were considered by Melrose and Uhlmann [25].

As applied to the seismic imaging problem, the results here have negative implications (as do the results of [28, 6] in the case of fold caustics): even microlocally away from $\Delta \cap \tilde{C}$, the non-pseudodifferential part of the normal operator has the same order as the pseudodifferential part, resulting in a strong, nonremovable artifact. However, our work provides a precise

description of the imaging artifacts resulting from cusp caustics, observed by Nolan and Symes [29] in a model 2D case. It would be of interest to obtain Sobolev estimates for operators associated with $\Delta \cup \widetilde{C}$, similar to the results in [8] for certain $I^{p,l}$ classes, including those arising from fold caustics.

We begin in Sec. 2 with a simple example from harmonic analysis of a FIO having two-sided cusps of the type that is relevant for the seismology problem, recall some basic singularity theory, and examine the corresponding normal operator. In Sec. 3, we describe the linearized inverse problem from seismology, and show that the canonical relation underlying the linearized forward scattering operator F is (i) associated with a two-sided cusp, but with the additional (highly nongeneric) features that (ii) the cusp points for the left and right projections are equal, and (iii) the images of the cusp points on both the left and right are involutive (coisotropic). In Sec. 4 we define a general class of canonical relations having this structure, and then derive a weak normal form, very close to the model examined in Sec. 2, for a general flat two-sided cusp. Finally, in Sec. 5, we use the oscillatory representations found in Sec. 4 to analyze the composition B^*A for two FIOs associated with such a canonical relation. We point out that the technique of deriving weak normal forms for degenerate FIOs and using them to prove composition theorems has its origins in [16] and was continued in [7]; different notions of weak normal forms were used to obtain estimates in [11, 12].

2 Example of an FIO with a flat two-sided cusp

2.1 A generalized Radon transform

We start with a simple example of a generalized Radon transform that incorporates the features of the linearized seismic inversion problem in which we are ultimately interested. In fact, we will later show that perturbations of this example serve as weak normal forms for the general class of operators with flat two-sided cusps.

The operator averaging over translates in $\mathbb{R}^4$ of a curve $\gamma(t)$ such that

$$\dot{\gamma}, \ddot{\gamma}, \dddot{\gamma}, \gamma^{(4)} \text{ are linearly independent} \quad (1)$$

is an FIO having a canonical relation which is a two-sided cusp [12], but the same holds for the curve $\gamma(t) = (t, t^2, t^4)$ in $\mathbb{R}^3$ [13, 14], and this is the model we will use.

Thus, consider the generalized Radon transform $R_0 : \mathcal{D}'(\mathbb{R}^3) \longrightarrow \mathcal{D}'(\mathbb{R}^3)$,

$$\begin{aligned} R_0 f(x) &= \int f(x - (t, t^2, t^4)) \chi(t) dt \\ &= \int_{\mathbb{R}^2} e^{i(x_2 - y_2 - (x_1 - y_1)^2)\theta_2 + (x_3 - y_3 - (x_1 - y_1)^4)\theta_3} f(y) \chi(x_1 - y_1) 1(\theta) d\theta_2 d\theta_3 dy, \end{aligned} \quad (2)$$

where $\chi \in C_0^\infty(\mathbb{R})$ is a fixed cutoff function. The associated canonical relation, the conormal bundle $C_0 = N^*Z' \subset (T^*\mathbb{R}^3 \setminus 0) \times (T^*\mathbb{R}^3 \setminus 0)$, where $Z = \text{supp}(K_{R_0}) \subset \mathbb{R}^3 \times \mathbb{R}^3$ and $(x, \xi; y, \eta)' = (x, \xi; y, -\eta)$ is standard notation for the twist map, is

$$\begin{aligned} C_0 = \Big\{ & \left(x_1, x_2, x_3, -2(x_1 - y_1)\theta_2 - 4(x_1 - y_1)^3\theta_3, \theta_2, \theta_3; \right. \\ & \left. y_1, y_2, y_3, -2(x_1 - y_1)\theta_2 - 4(x_1 - y_1)^3\theta_3, \theta_2, \theta_3 \right) : \\ & \left. x_2 - y_2 - (x_1 - y_1)^2 = x_3 - y_3 - (x_1 - y_1)^4 = 0 \right\}. \end{aligned} \quad (3)$$

We will show that C_0 has several properties, which remarkably also hold in the totally unrelated seismic imaging problem: (i) the projections both to the left and right, $\pi_L, \pi_R : C_0 \longrightarrow T^*\mathbb{R}^3 \setminus 0$, have (Whitney) cusp degeneracies; (ii) the cusp points for the two projections are the same, $\Sigma_{1,1}(\pi_L) = \Sigma_{1,1}(\pi_R) := \Sigma_{1,1}$; and (iii) the images of the cusp points, $\pi_L(\Sigma_{1,1})$ and $\pi_R(\Sigma_{1,1})$, are coisotropic (involutive) submanifolds of $T^*\mathbb{R}^3$. Conditions (ii) and (iii) are unstable and quite special among two-sided cusps, i.e., canonical relations satisfying (i); see the Remarks in Sec. 2.5 below.

2.2 Singularity classes

We first recall some basic facts about cusps and refer to [33, 27, 10] for more details. Let $f : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be a smooth function. We say that f *drops rank simply* at p if $\text{rank}(df)_p = N - 1$ and if $(d(\det df))_p \neq 0$, so that $\Sigma_1(f) := \{x \in \mathbb{R}^N : \det(df(x)) = 0\}$, the corank one critical set of f , is a smooth hypersurface near p . If $\ker df_p \not\subset T_p \Sigma_1(f)$, then f has a *fold* singularity, and $f|_{\Sigma_1(f)}$ is an immersion. Considering the more degenerate case when $\ker df_p \subset T_p \Sigma_1(f)$, one may choose a nonzero vector field v along

$\Sigma_1(f)$ such that $v \in \ker df_p$, so that v is tangent to $\Sigma_1(f)$ at p . Let g be a smooth function such that $g|_{\Sigma_1(f)} = 0$ and $dg_p \neq 0$, e.g., $g = \det df$. Thus, $dg(v)$ has a zero at p .

Definition 2.1. f has a (Whitney) *cuspidal set* at p if $dg(v)$ has a simple zero at p .

$\Sigma_{1,1}(f)$, the *cuspidal set* of f , is then codimension 2, and $f|_{\Sigma_{1,1}(f)}$ is an immersion. One may use adapted coordinates to clarify this. These are local coordinates such that $f(x_1, x_2, \dots, x_N) = (x_1, x_2, \dots, x_{N-1}, h(x))$ and $h(0) = 0$ [27]. Then, $\Sigma_1(f) = \{x : \frac{\partial h}{\partial x_N}(0) = 0\}$, and f has a cusp singularity at 0 iff $\frac{\partial^2 h}{\partial x_N^2}(0) = 0$, $\frac{\partial^3 h}{\partial x_N^3}(0) \neq 0$ and $\text{rank} [d_x(\frac{\partial h}{\partial x_N}), d_x(\frac{\partial^2 h}{\partial x_N^2})] = 2$. These conditions are adapted coordinate-independent, and the notion of a cusp makes sense for any smooth mapping between N -dimensional manifolds. In suitable smooth coordinates on the domain and range spaces, any map with a cusp singularity can be put into the local normal form, $f(x_1, x_2, \dots, x_N) = (x_1, x_2, \dots, x_{N-1}, x_{N-1}x_N + x_N^3)$.

For the canonical relation C_0 , the projection to the left, $\pi_L : C_0 \rightarrow T^*\mathbb{R}^3$, is $\pi_L(x_1, x_2, x_3, y_1, \theta_2, \theta_3) = (x_1, x_2, x_3, -2(x_1 - y_1)\theta_2 - 4(x_1 - y_1)^3\theta_3, \theta_2, \theta_3)$; hence, letting $\alpha = 2\theta_2 + 12(x_1 - y_1)^2\theta_3$, $\beta = -2(x_1 - y_1)$ and $\gamma = -4(x_1 - y_1)^3$, one has

$$d\pi_L = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ -\alpha & 0 & 0 & \alpha & \beta & \gamma \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

which has $\det d\pi_L = 2\theta_2 + 12(x_1 - y_1)^2\theta_3$ and $\ker \pi_L = \mathbb{R} \cdot \frac{\partial}{\partial y_1}$ at $\Sigma_1(\pi_L) = \{2\theta_2 + 12(x_1 - y_1)^2\theta_3 = 0\}$. Since $\frac{\partial}{\partial y_1}(2\theta_2 + 12(x_1 - y_1)^2\theta_3) = -24(x_1 - y_1)\theta_3$ and $\frac{\partial^2}{\partial y_1^2}(2\theta_2 + 12(x_1 - y_1)^2\theta_3) = 24\theta_3 \neq 0$, the cuspidal set is

$$\Sigma_{1,1}(\pi_L) = \{2\theta_2 + 12(x_1 - y_1)^2\theta_3 = x_1 - y_1 = 0\} = \{\theta_2 = x_1 - y_1 = 0\}.$$

Noting that $2\theta_2 + 12(x_1 - y_1)^2\theta_3$ and $x_1 - y_1$ have linearly independent gradients, we see that π_L has a cusp singularity.

Similarly, the projection to the right is

$$\pi_R(x, y_1, \theta_2, \theta_3) = (y_1, x_2 - (x_1 - y_1)^2, x_3 - (x_1 - y_1)^4, -2(x_1 - y_1)\theta_2 - 4(x_1 - y_1)^3\theta_3, \theta_2, \theta_3),$$

so that $\det d\pi_R = -2\theta_2 - 12(x_1 - y_1)^2\theta_3$, and $\ker \pi_R = \frac{\partial}{\partial x_1}$ at $\Sigma_1(\pi_R) = \{2\theta_2 + 12(x_1 - y_1)^2\theta_3 = 0\}$. One also has $\frac{\partial}{\partial x_1}(2\theta_2 + 12(x_1 - y_1)^2\theta_3) = 24(x_1 - y_1)\theta_3$ and $\frac{\partial^2}{\partial x_1^2}(2\theta_2 + 12(x_1 - y_1)^2\theta_3) = 24\theta_3 \neq 0$, so that $\Sigma_{1,1}(\pi_R) = \{\theta_2 = x_1 - y_1 = 0\}$, and thus π_R also has a cusp singularity.

Note that $\Sigma_1(\pi_R) = \Sigma_1(\pi_L)$ and $\Sigma_{1,1}(\pi_R) = \Sigma_{1,1}(\pi_L) =: \Sigma_{1,1}$. The first of these is true for any canonical relation [19], but the second is a very strong condition; indeed, for general two-sided cusps, there is no relationship between $\Sigma_{1,1}(\pi_L)$ and $\Sigma_{1,1}(\pi_R)$. Furthermore, for the images of these sets, one has

$$\pi_L(\Sigma_1) = \{\xi_2^3 = -\frac{27}{8}\xi_1^2\xi_3\}, \quad \pi_R(\Sigma_1) = \{\eta_2^3 = -\frac{27}{8}\eta_1^2\eta_3\},$$

with $\pi_L(\Sigma_{1,1}) = \{\xi_1 = \xi_2 = 0\}$, $\pi_R(\Sigma_{1,1}) = \{\eta_1 = \eta_2 = 0\}$, resp., their cuspidal edges. $\pi_L(\Sigma_{1,1})$ and $\pi_R(\Sigma_{1,1})$ are codimension two coisotropic (or involutive) submanifolds of $T^*\mathbb{R}^3 \setminus 0$ [19]; again, this is a nongeneric situation.

Consider next the composition of canonical relations,

$$C_0^t \circ C_0 = \{(x, \xi; y, \eta) : \exists(z, \zeta) \text{ s.t. } (x, \xi; z, \zeta) \in C_0^t \text{ and } (z, \zeta; y, \eta) \in C_0\}.$$

For such a triple $(x, \xi; z, \zeta; y, \eta)$, one obtains the following equations:

$$\begin{aligned} z_2 - x_2 &= (z_1 - x_1)^2, & z_3 - x_3 &= (z_1 - x_1)^4, \\ \xi_1 = \zeta_1 &= -2(z_1 - x_1)\zeta_2 - 4(z_1 - x_1)^3\zeta_3, & \xi_2 = \zeta_2, & \xi_3 = \zeta_3, \\ z_2 - y_2 &= (z_1 - y_1)^2, & z_3 - y_3 &= (z_1 - y_1)^4, \\ \eta_1 = \zeta_1 &= -2(z_1 - y_1)\zeta_2 - 4(z_1 - y_1)^3\zeta_3, & \eta_2 = \zeta_2, & \eta_3 = \zeta_3. \end{aligned}$$

Using $2(z_1 - x_1)\xi_2 + 4(z_1 - x_1)^3\xi_3 = 2(z_1 - y_1)\xi_2 + 4(z_1 - y_1)^3\xi_3$, after simplification one obtains:

$$\begin{aligned} (y_1 - x_1)\xi_2 + 2[(z_1 - x_1)^3 - (z_1 - y_1)^3]\xi_3 &= 0, \\ (y_1 - x_1)[\xi_2 + 2(3z_1^2 - 3z_1(x_1 + y_1) + x_1^2 + y_1^2 + x_1y_1)\xi_3] &= 0 \end{aligned}$$

It follows that the contribution to $C_0^t \circ C_0$ from $\{y_1 - x_1 = 0\}$ is contained in Δ , and that from $\{y_1 - x_1 \neq 0\}$ is contained in $\tilde{C}_0$, where

$$\begin{aligned} \tilde{C}_0 = \Big\{ & (x_1, x_2, x_3, -2(z_1 - x_1)\theta_2 - 4(z_1 - x_1)^3\theta_3, \theta_2, \theta_3; \\ & y_1, y_2, y_3, -2(z_1 - x_1)\theta_2 - 4(z_1 - x_1)^3\theta_3, \theta_2, \theta_3) : \\ & x \in \mathbb{R}^3, y_1, z_1 \in \mathbb{R}, \theta_3 \in \mathbb{R} \setminus 0, y_2 = x_2 + (y_1 - x_1)(2z_1 - x_1 - y_1), \\ & y_3 = x_3 + (y_1 - x_1)(2z_1 - x_1 - y_1)((z_1 - x_1)^2 + (z_1 - y_1)^2), \\ & \theta_2 = -2(3z_1^2 - 3z_1(x_1 + y_1) + x_1^2 + y_1^2 + x_1y_1)\theta_3 \Big\} \end{aligned} \quad (4)$$

Notice that $\xi_1 = \eta_1 = 4\theta_3(z_1 - x_1)(z_1 - y_1)(2z_1 - x_1 - y_1)$ and that $\tilde{C}_0$ intersects Δ in codimension one, at $\{x_1 - y_1 = 0\}$.

The parametrization of $\tilde{C}_0$ in (4) is a map

$$\Upsilon : \mathbb{R}_{x_1, x_2, x_3, y_1, z_1}^5 \times (\mathbb{R}_{\theta_3} \setminus 0) \longrightarrow (T^*\mathbb{R}^3 \setminus 0) \times (T^*\mathbb{R}^3 \setminus 0).$$

One easily sees that Υ is singular at $\tilde{\Sigma} := \Sigma_1(\Upsilon) = \{x_1 - y_1 = x_1 - z_1 = 0\}$, where $\ker d\Upsilon = \mathbb{R} \cdot \frac{\partial}{\partial z_1}$, which is $\not\subset T\Sigma_1$. Thus, as discussed below, $\tilde{C}_0$ is an *open umbrella*, exhibiting the simplest kind of singularity of a Lagrangian manifold.

2.3 Open umbrellas

For maps between manifolds of the same dimension, a fold is a singularity of the type $S_{1,0}$ and a cusp is a singularity of type $S_{1,1,0}$ [33, 27, 10]. From larger dimensional spaces to smaller ones, $S_{1,0}$ maps are submersions with folds, while in the opposite direction they are referred to variously as *cross caps* or *Whitney-Cayley umbrellas* [10, 9]. In the lowest possible dimensions, if $g : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ has a cross cap singularity then, in suitable local coordinates, $(u, v, w) = g(x, y) = (x^2, y, xy)$, and its image is the algebraic surface $\{w^2 = uv^2\}$, the Whitney-Cayley umbrella. This is the simplest type of non-immersed surface singularity in three dimensions. It is an immersion away from the origin; it is actually an embedding on $\{y \neq 0\}$; and along $\{y = 0\}$, it is folded and hence 2-1.

Now, by adding one dimension to the range space, one may simultaneously both unfold the closed umbrella, making the parametrization 1-1 away from the origin, and make it a Lagrangian manifold with singularity in

$\mathbb{R}^4$. The *open* (or unfolded) *umbrella* is the map $U : R^2 \rightarrow R^4$, $U(x, y) = (x^2, y, xy, \frac{2}{3}x^3)$ (and its image). We have $U^*\omega = 0$, where ω is the symplectic form on $R^4 \simeq T^*\mathbb{R}^2$. Hence, the unfolded umbrella is a *Lagrangian inclusion*, i.e., a variety which is a smoothly immersed Lagrangian manifold away from its singular points [9]. To find a similar conic model, one needs to look in $T^*R^3 \setminus 0$; for an unknown function $f(t, s)$, consider

$$\Lambda_1 = \left\{ (s^2, t, f(t, s); st\theta, \frac{2}{3}t^3\theta, \theta) \in T^*\mathbb{R}^3 \setminus 0 : \theta \neq 0 \right\}.$$

Then, Λ_1 is a Lagrangian if f is chosen so that $d\xi \wedge dx|_{\Lambda_1} = 0$. Since $d\xi \wedge dx = d(st\theta) \wedge d(t^2) + d(\frac{2}{3}t^3\theta) \wedge ds + d\theta \wedge df = d\theta \wedge (2st^2dt + \frac{2}{3}t^3ds) + d\theta \wedge df$, this holds if $df = -2st^2dt - \frac{2}{3}t^3ds = -d(\frac{2}{3}t^3s)$; using $f = -\frac{2}{3}t^3s$ yields the conic Lagrangian,

$$\Lambda_1 = \left\{ (t^2, s, -\frac{2}{3}t^3s; st\theta, \frac{2}{3}t^3\theta, \theta) : t, s \in \mathbb{R}, \theta \in \mathbb{R} \setminus 0 \right\},$$

exhibiting an open umbrella singularity along $\{(0, 0, 0; 0, 0, \xi_3) : \xi_3 \neq 0\}$.

Observe that Λ_1 may be parametrized by a degenerate phase function. Letting $\varphi(x, \theta_1, \theta_2, \tau, \eta) = (x_1 - (\frac{\tau}{\eta})^2)\theta_1 + (x_3 + \frac{2}{3}(\frac{\tau}{\eta})^3x_2)\theta_2$, one has

$$d_{\theta_1}\varphi = x_1 - (\frac{\tau}{\eta})^2, \quad d_{\theta_2}\varphi = x_3 + \frac{2}{3}(\frac{\tau}{\eta})^3x_2$$

$$d_{\tau}\varphi = \frac{\tau}{\eta}(-2\frac{\theta_1}{\eta} + 2\frac{\tau}{\eta}\frac{\theta_2}{\eta}x_2), \quad d_{\eta}\varphi = -(\frac{\tau}{\eta})^2(-2\frac{\theta_1}{\eta} + 2\frac{\tau}{\eta}\frac{\theta_2}{\eta}x_2),$$

so that φ is degenerate, with both $d_{\tau}\varphi$ and $d_{\eta}\varphi$ having normal crossings: $d_{\tau}\varphi = 0$ iff $\tau = 0$ or $-2\frac{\theta_1}{\eta} + 2\frac{\tau}{\eta}\frac{\theta_2}{\eta}x_2 = 0$, and $d_{\eta}\varphi = 0$ on same two sets. For the second case, solving for $\theta_1 = \frac{\tau}{\eta}x_2\theta_2$ we see that φ parametrizes Λ_1 , while for the first case, φ also parametrizes $\Lambda_0 := N^*\{x_1 = x_3 = 0\}$. Λ_0 and Λ_1 intersect cleanly in codimension one, except at the singular set of Λ_1 , which is contained in the intersection. φ simultaneously parametrizes $\Lambda_0 \cup \Lambda_1$, although it is not a multiphase function in the sense of [26], used to parametrize a pair of cleanly intersecting smooth Lagrangians. The class of generalized FIOs we define in Sec. 5.1 has similar features.

For the general notion of an open umbrella, let (M, ω) be a smooth symplectic manifold of dimension $2n$.

Definition 2.2. $S \subset M$ is an *open umbrella* if it is the image of a map $\psi : R^n \rightarrow M$ such that $d\psi$ drops rank by one simply at Σ_1 , of codim two; $\ker (d\psi)|_{\Sigma_1} \not\subset T\Sigma_1$; and $\text{Range} (d\psi_p)$ is Lagrangian for $p \in R^n \setminus \Sigma$.

The class of open umbrellas is structurally stable under smooth perturbations in the class of Lagrangian inclusions [9]. As checked above, the parametrization Υ in (4) satisfies the conditions in the definition, so that $\tilde{C}_0$ is an open umbrella.

2.4 Composition for the model flat two sided cusp

We now want to show that the composition A^*A for Fourier integral operators, such as R_0 , associated with the model flat two-sided cusp C_0 in (3) results in operators with wave front relation in the union of the diagonal Δ and the open umbrella $\tilde{C}_0$ from (4). Let $A \in I^m(\mathbb{R}^3, \mathbb{R}^3; C_0)$, so that $Af(x) = \int e^{i\phi(x,y,\theta)} a(x,y,\theta) f(y) d\theta dy$ with $\phi(x,y,\theta) = (x_2 - y_2 - (x_1 - y_1)^2)\theta_2 + (x_3 - y_3 - (x_1 - y_1)^4)\theta_3$ and $a \in S_{1,0}^{m+\frac{1}{2}}$, the standard Hörmander class of symbols. We consider the normal operator formed by the composition,

$$A^*Af(x) = \int e^{i(\phi(z,y,\eta) - \phi(z,x,\theta))} a(z,y,\eta) \bar{a}(z,x,\theta) f(y) dy dz d\theta d\eta.$$

We have

$$\begin{aligned} \phi(z,y,\eta) - \phi(z,x,\theta) &= (z_2 - y_2 - (z_1 - y_1)^2)\eta_2 + (z_3 - y_3 - (z_1 - y_1)^4)\eta_3 \\ &\quad - (z_2 - x_2 - (z_1 - x_1)^2)\theta_2 - (z_3 - x_3 - (z_1 - x_1)^4)\theta_3. \end{aligned}$$

After a stationary phase in (z_2, η_2) and (z_3, η_3) in the integral for K_{A^*A} , we obtain:

$$\begin{aligned} \phi(z,y,\eta) - \phi(z,x,\theta) &= (x_2 - y_2 + (z_1 - x_1)^2 - (z_1 - y_1)^2)\theta_2 + (x_3 - y_3 + (z_1 - x_1)^4 - (z_1 - y_1)^4)\theta_3 \\ &= (x_2 - y_2)\theta_2 + (x_3 - y_3)\theta_3 + (y_1 - x_1)[(2z_1 - x_1 - y_1)\theta_2 + (2z_1 - y_1 - x_1)((z_1 - x_1)^2 + (z_1 - y_1)^2)\theta_3] \end{aligned}$$

Let $z_1 = \frac{\zeta}{\theta_3}$, and then consider the change of variable,

$$\theta_1 = -(2\frac{\zeta}{\theta_3} - x_1 - y_1)\theta_2 - (2\frac{\zeta}{\theta_3} - y_1 - x_1)((\frac{\zeta}{\theta_3} - x_1)^2 + (\frac{\zeta}{\theta_3} - y_1)^2)\theta_3,$$

for which

$$\begin{aligned}\frac{\partial \theta_1}{\partial \zeta} &= -2\frac{\theta_2}{\theta_3} - 2\left(\left(\frac{\zeta}{\theta_3} - x_1\right)^2 + \left(\frac{\zeta}{\theta_3} - y_1\right)^2\right) - 2\left(2\frac{\zeta}{\theta_3} - x_1 - y_1\right)^2, \\ \frac{\partial^2 \theta_1}{\partial \zeta^2} &= -\frac{12}{\theta_3}\left(2\frac{\zeta}{\theta_3} - x_1 - y_1\right), \text{ and } \frac{\partial^3 \theta_1}{\partial \zeta^3} = -\frac{24}{\theta_3}.\end{aligned}$$

When $\frac{\partial^2 \theta_1}{\partial \zeta^2} = 0$ then $2\frac{\zeta}{\theta_3} = x_1 + y_1$, $\theta_1 = 0$ and $\theta_2 = -\frac{(x_1 - y_1)^2}{2}\theta_3$. Let $T(t_1, t_2) = (t_1 t_2 + \frac{1}{2}t_1^3, t_2)$ be a normal form for a two-dimensional map with a cusp singularity. To understand the push forward T_*u of a distribution u under the map T , note that the kernel of T_* is

$$K_{T_*}(z, t) = \delta(z - T(t)) = \int e^{i[(z_1 - t_1 t_2 - \frac{1}{2}t_1^3)\sigma_1 + (z_2 - t_2)\sigma_2]} 1(\sigma) d\sigma,$$

so that

$$T_*u(z) = \int e^{i[z_1\sigma_1 + z_2\sigma_2]} \left[\int e^{-i[t_2\sigma_2 + (t_1 t_2 + \frac{1}{2}t_1^3)\sigma_1]} u(t) dt \right] d\sigma.$$

Returning now to A^*A , letting $z_1 = \frac{\theta_1}{\theta_3}$ and $z_2 = \frac{\theta_2}{\theta_3} + \frac{1}{2}(x_1 - y_1)^2$, the phase function of A^*A becomes

$$\begin{aligned}\tilde{\phi}(x, y, \theta, \sigma) &= (x_1 - y_1)\theta_1 + (x_2 - y_2)\theta_2 + (x_3 - y_3)\theta_3 \\ &\quad + \left(\frac{\theta_1}{\theta_3} - t_1 t_2 - \frac{1}{2}t_1^3\right)\sigma_1 + \left(\frac{\theta_2}{\theta_3} + \frac{1}{2}(x_1 - y_1)^2 - t_2\right)\sigma_2\end{aligned}$$

and the amplitude becomes: $a \times 1(\sigma)$. Next, we perform stationary phase, first in t_2, σ_2 and then in θ_1, σ_1 . We have

$$d_{t_2}\tilde{\phi} = -t_1\sigma_1 - \sigma_2, \quad d_{\sigma_2}\tilde{\phi} = \frac{\theta_2}{\theta_3} + \frac{1}{2}(x_1 - y_1)^2 - t_2.$$

The Hessian is 1, the phase function is $\tilde{\phi} = (x - y)\theta + (\frac{\theta_1}{\theta_3} - \frac{1}{2}t_1^3 - t_1[\frac{\theta_2}{\theta_3} + \frac{1}{2}(x_1 - y_1)^2])\sigma_1$, and

$$d_{\theta_1}\tilde{\phi} = x_1 - y_1 + \frac{\sigma_1}{\theta_3}, \quad d_{\sigma_1}\tilde{\phi} = \frac{\theta_1}{\theta_3} - \frac{1}{2}t_1^3 - t_1\left(\frac{\theta_2}{\theta_3} + \frac{1}{2}(x_1 - y_1)^2\right).$$

The Hessian is $\frac{1}{\theta_3^2}$. Finally, we introduce the variable $\tau = t_1\theta_3$ and obtain

$$\tilde{\phi} = \left(x_2 - y_2 + \frac{\tau}{\theta_3}(x_1 - y_1)\right)\theta_2 + \left(x_3 - y_3 + \frac{1}{2}(x_1 - y_1)\left(\frac{\tau}{\theta_3}\right)^3 + \frac{1}{2}\frac{\tau}{\theta_3}(x_1 - y_1)^3\right)\theta_3,$$

with respect to which the Schwartz kernel has the representation

$$K_{A^*A}(x, y) = \int e^{i\tilde{\phi}} b \, d\theta_2 d\theta_3 d\tau, \quad b \in S^{2m}.$$

The wave front relation satisfies

$$\begin{aligned} WF(K_{A^*A})' \subset & \left\{ (x_1, x_2, x_3, \frac{\tau\theta_2}{\theta_3} + \frac{1}{2}\frac{\tau^3}{\theta_3^2} + \frac{3}{2}(x_1 - y_1)^2\tau, \theta_2, \theta_3; \right. \\ & y_1, y_2, y_3, \frac{\tau\theta_2}{\theta_3} + \frac{1}{2}\frac{\tau^3}{\theta_3^2} + \frac{3}{2}(x_1 - y_1)^2\tau, \theta_2, \theta_3) : \\ & d_{\theta_2}\tilde{\phi} = x_2 - y_2 + \frac{\tau}{\theta_3}(x_1 - y_1) = 0, \\ & d_{\theta_3}\tilde{\phi} = x_3 - y_3 - \frac{\theta_2\tau}{\theta_3^2}(x_1 - y_1) - \frac{\tau^3}{\theta_3^3}(x_1 - y_1) = 0, \\ & \left. d_{\tau}\tilde{\phi} = \frac{\theta_2}{\theta_3}(x_1 - y_1) + \frac{3}{2}\frac{\tau^2}{\theta_3^2}(x_1 - y_1) + \frac{1}{2}(x_1 - y_1)^3 = 0 \right\}. \end{aligned}$$

From the last relation, we see that the phase function is degenerate, with the critical set in the phase variables having a normal crossing, being the union of two transverse surfaces, corresponding to $x_1 - y_1 = 0$ or $\frac{\theta_2}{\theta_3} + \frac{3}{2}\frac{\tau^2}{\theta_3^2} + \frac{1}{2}(x_1 - y_1)^2 = 0$. The points where $x_1 = y_1$ give rise to the diagonal Δ , but with the parametrization actually being a cusp:

$$(x, \tau, \theta) \longrightarrow \left(x, \frac{\tau\theta_2}{\theta_3} + \frac{1}{2}\frac{\tau^3}{\theta_3^2}, \theta_2, \theta_3; \quad x, \frac{\tau\theta_2}{\theta_3} + \frac{1}{2}\frac{\tau^3}{\theta_3^2}, \theta_2, \theta_3 \right).$$

On the other hand, the points where $\frac{\theta_2}{\theta_3} + \frac{3}{2}\frac{\tau^2}{\theta_3^2} + \frac{1}{2}(x_1 - y_1)^2 = 0$ contribute to $WF(A^*A)$ the set, parametrized by $x \in \mathbb{R}^3, y_1, \tau \in \mathbb{R}, \theta_3 \neq 0$,

$$\begin{aligned} \tilde{C}_0 = \left\{ \begin{aligned} (x, \xi; y, \eta) : y_2 &= x_2 + \frac{\tau}{\theta_3}(x_1 - y_1), & \xi_3 &= \eta_3 = \theta_3, \\ y_3 &= x_3 + \frac{1}{2}\frac{\tau}{\theta_3}(x_1 - y_1)^3 + \frac{1}{2}\frac{\tau^3}{\theta_3^3}(x_1 - y_1), \\ \xi_1 &= \eta_1 = -\frac{\tau^3}{\theta_3^3} + (x_1 - y_1)^2\tau, \\ \xi_2 &= \eta_2 = -\frac{1}{2}(x_1 - y_1)^2\theta_3 - \frac{3}{2}\frac{\tau^2}{\theta_3}, \end{aligned} \right\}, \end{aligned} \quad (5)$$

which is the image of $\rho(x_1, x_2, x_3, y_1, \theta_3, \tau) = (x, \xi; y, \eta)$ with $y_2, y_3, \eta_1, \eta_2, \eta_3, \xi_1, \xi_2, \xi_3$ given by the above relations. ρ satisfies the conditions of Def. 2.2: $d\rho$ drops rank simply by one at $\Sigma = \{x_1 - y_1 = \tau = 0\}$ and $\ker d\rho = \mathbb{R} \cdot \frac{\partial}{\partial \tau} \notin T\Sigma$. Hence $\tilde{C}_0$ is an open umbrella.

Next, we compute the principal symbols on Δ and $\tilde{C}_0$ away from their intersection. We have $\text{Crit}_{\tilde{\phi}} = \text{Crit}_{\Delta} \cup \text{Crit}_{\tilde{C}_0} = \{(x, y, \tau, \theta_2, \theta_3) \mid d_{\tau}\tilde{\phi} = 0, d_{\theta_2}\tilde{\phi} = 0, d_{\theta_3}\tilde{\phi} = 0\}$. Using Hörmander's formula [19], $\sigma = a(E_{\tilde{\phi}})^{\frac{1}{2}}$ where $E_{\tilde{\phi}} = |\frac{D(\lambda_i, \frac{\partial \tilde{\phi}}{\partial \theta})}{D(x, \theta)}|^{-1}$ and λ_i are local coordinates on $\text{Crit}_{\tilde{\phi}}$. On Crit_{Δ} , local coordinates are $(x, \tau, \theta_2, \theta_3)$ and $E_{\tilde{\phi}} = |\frac{D(x, \tau, \theta_2, \theta_3, \frac{\partial \tilde{\phi}}{\partial \tau}, \frac{\partial \tilde{\phi}}{\partial \theta_2}, \frac{\partial \tilde{\phi}}{\partial \theta_3})}{D(x, \tau, \theta_2, \theta_3, y_1, y_2, y_3)}|^{-1} = (\frac{\theta_2}{\theta_3} + \frac{3}{2} \frac{\tau^2}{\theta_3^2})^{-1}$, while, on $\text{Crit}_{\tilde{C}_0}$, local coordinates are (x, y_1, τ, θ_3) and $E_{\tilde{\phi}} = |\frac{D(x, y_1, \tau, \theta_3, \frac{\partial \tilde{\phi}}{\partial \tau}, \frac{\partial \tilde{\phi}}{\partial \theta_2}, \frac{\partial \tilde{\phi}}{\partial \theta_3})}{D(x, y_1, \tau, \theta_2, \theta_3, y_2, y_3, \theta_2)}|^{-1} = (\frac{x_1 - y_1}{\theta_3})^{-1}$. Thus, both principal symbols are singular as one approaches $\Delta \cap \tilde{C}_0$, behaving as $\delta^{-\frac{1}{2}}$, where δ is the distance on either Δ or $\tilde{C}_0$ to the intersection.

2.5 Flat two-sided cusps

Motivated by the previous example, we now define the class of canonical relations for which we will establish a composition calculus.

Definition 2.3. If X and Y are manifolds of dimension $n \geq 3$, then a canonical relation $C \subset (T^*X \setminus 0) \times (T^*Y \setminus 0)$ is a *flat two-sided cusp* if

- (i) both $\pi_L : C \rightarrow T^*X$ and $\pi_R : C \rightarrow T^*Y$ have at most cusp singularities;
- (ii) the left- and right-cusp points are equal: $\Sigma_{1,1}(\pi_L) = \Sigma_{1,1}(\pi_R) =: \Sigma_{1,1}$; and
- (iii) $\pi_L(\Sigma_{1,1}) \subset T^*X$ and $\pi_R(\Sigma_{1,1}) \subset T^*Y$ are coisotropic (involutive) and nonradial.

Here, as usual, *at most cusp* means either cusps or folds, which are unavoidable in the neighborhood of a cusp. *Nonradial*, meaning that the radial vector field $\sum \xi_j \partial_{\xi_j}$ does not lie in the two-dimensional Hamiltonian foliation of the codimension two coisotropic submanifold, is a standard technical assumption needed to apply the homogeneous Darboux theorem in Sec. 4.

Other examples of flat two-sided cusps coming from generalized Radon transforms, verified as with the example above, are those for translates of

curves in $\mathbb{R}^4$ satisfying (1), and for the translates in $\mathbb{R}^3$ of Menn's surface [1],

$$C_1 = N^*\{x_3 - y_3 = (x_1 - y_1)^2(x_2 - y_2) - (x_2 - y_2)^2\}',$$

or any other surface in $\mathbb{R}^3$ whose Gauss map has a cusp. Computation also shows that the composition $C_1^t \circ C_1 = \Delta \cup \tilde{C}_1$, with $\tilde{C}_1$ similar to $\tilde{C}_0$ as in (5) above.

Remarks. 1. If (i) and (ii) in Def. 2.3 are satisfied, the structure of $\omega_{T^*X}|_{T\pi_L(\Sigma_{1,1})}$ is the same as $\omega_{T^*Y}|_{T\pi_R(\Sigma_{1,1})}$, since both are equivalent to the restriction to $\Sigma_{1,1} \subset C$ of the degenerate symplectic form $\omega_C := \pi_L^* \omega_{T^*X} = \pi_R^* \omega_{T^*Y}$. Thus, $\pi_L(\Sigma_{1,1})$ is involutive iff $\pi_R(\Sigma_{1,1})$ is, and one can see that this holds iff the two-dimensional $\ker(\omega_C|_{\Sigma_1})$ is simply tangent to Σ_1 at $\Sigma_{1,1}$. This is in contrast to the situation when the image of the cusp set is symplectic (strongly noninvolutive), in which case ω_C is a folded symplectic form and $\ker(\omega_C|_{\Sigma_1}) \pitchfork \Sigma_1$ everywhere [22].

2. Condition (i) is stable under small perturbations, since cusps are structurally stable [10]. However, given (i), condition (ii) is unstable and atypical. To see this concretely, consider (possibly) nontranslationally invariant families of curves $\{\gamma_x\}_{x \in \mathbb{R}^4}$, using the framework of [4] as analyzed in terms of FIOs with cusps in [12]. For vector fields X, Y, Z, W on $\mathbb{R}^4$, with $X \neq 0$, let

$$\gamma_x(t) = \exp_x(tX + t^2Y + t^3Z + t^4W).$$

The associated generalized Radon transform is in $I^{-\frac{1}{2}}(\mathbb{R}^4, \mathbb{R}^4; C)$, with C a canonical relation for which, by [12, Prop. 6.1], π_L (resp., π_R) is at most a cusp if

$$X, Y, Z \pm \frac{1}{6}[X, Y], W \pm \frac{1}{4}[X, Z] + \frac{1}{24}[X, [X, Y]] \text{ are linearly independent.}$$

From the calculations in [12], one sees that a necessary condition for $\Sigma_{1,1}(\pi_L) = \Sigma_{1,1}(\pi_R)$ is that $(Z + \frac{1}{6}[X, Y]) - (Z - \frac{1}{6}[X, Y]) \in \text{span}\{X, Y\}$, i.e., $\text{span}\{X, Y\}$ is an integrable distribution of 2-planes, which is generically not the case.

Nevertheless, as seen in the next section, flat two-sided cusps arise naturally in a quite different setting without translation invariance or integrability.

3 Linearized seismic inversion with cusp caustics

We consider linearized seismic inversion for the single source data set, under the assumption that the background sound speed has caustics of at most cusp type, i.e., either folds or cusps. We show that the linearized forward scattering operator F is a Fourier integral operator associated with a flat two-sided cusp, $\tilde{C}$. We only briefly describe this problem, referring to [29, 30, 28, 6, 7] for more motivation and details.

Acoustic waves are generated at the surface of the earth, scatter off heterogeneities in the subsurface and return to the surface, where measurements of the pressure field are used to reconstruct an image of the subsurface. The model for the scattered waves is given by the acoustic wave equation,

$$\frac{1}{c^2(x)} \frac{\partial^2 p}{\partial t^2}(x, t) - \Delta p(x, t) = \delta(x - s) \delta(t), \quad p(x, t) = 0, \quad t < 0, \quad (6)$$

where $x \in Y = \{x \in \mathbb{R}^3, x_3 \geq 0\}$, representing the Earth, $p(x, t)$ is the pressure field resulting from a pulse at the source s at time $t = 0$, and $c(x)$ is the unknown sound speed field.

To make this nonlinear inverse problem tractable, one considers a linearized operator F which maps singular perturbations of a smooth background sound speed in the subsurface, assumed known, to perturbations of the resulting pressure field at the surface. Thus, the linearization consists in assuming c to be of the form $c = c_0 + \delta c$ and the resulting $p = p_0 + \delta p$, where c_0 is a smooth known background field. The formal linearization of (6) is

$$\square_{c_0}(\delta p) := \frac{1}{c_0^2(x)} \frac{\partial^2 \delta p}{\partial t^2}(x, t) - \Delta \delta p(x, t) = \frac{2}{c_0^3} \frac{\partial^2 p_0}{\partial t^2} \cdot \delta c(x), \quad \delta p = 0, \quad t < 0, \quad (7)$$

where p_0 , the Green's function for $\square_{c_0}$. The *linearized scattering operator* is $F : \delta c \rightarrow \delta p|_{\partial Y \times (0, T)}$. Under mild technical assumptions, F is an FIO associated with a canonical relation $C \subset (T^*X \setminus 0) \times (T^*Y \setminus 0)$, where $X = \partial Y \times (0, T)$ is the data space [31, 20, 30]. The goal is then to left-invert F ; standard techniques suggest studying left invertibility of the normal operator, $N = F^*F$.

Let $H(x, \xi) = \frac{1}{2}(c_0(x)^{-2} - |\xi|^2)$ be the Hamiltonian associated to c_0 , and Λ_s the image of $T_s^*\mathbb{R}^3 \setminus 0$ under the bicharacteristic flow associated to H , which is a Lagrangian submanifold of $T^*\mathbb{R}^3 \setminus 0$. A *caustic* is a singularity of

the spatial projection $\pi : \Lambda_s \rightarrow Y$. It is known that the caustics exhibited for generic soundspeeds c_0 are the same as generic Lagrangian singularities, i.e., folds, cusps, swallowtails, etc. [17].

If c_0 is such that only fold caustics occur, it was shown by Nolan [28] that $F \in I^1(X, Y; C)$ with C a two-sided fold. Furthermore, the Schwartz kernel of the operator F^*F belongs to a class of distributions associated to two cleanly intersecting Lagrangians in $(T^*Y \setminus 0) \times (T^*Y \setminus 0)$, with the corresponding canonical relations being the diagonal Δ and another folding canonical relation [28, 6].

The next caustics to consider are those of cusp type, meaning that the only singularities of the spatial projection $\pi : \Lambda_s \rightarrow Y$ are folds or cusps. A cusp caustic is already present in the 2D example of [29] exhibiting loss of regularity for F . To analyze the geometry of the canonical relation C in the presence of cusp caustics, we make use of the description of Λ_s in [28]. It can be parametrized by t_{inc} , the time travelled by the incident ray, and the takeoff direction $(p_1, p_2, p_3) \in \mathbb{S}^2$. We can change these coordinates to (x_1, x_2, p_3) . Hence on $\tilde{C}_s$, $x_3 = f(x_1, x_2, p_3)$ and $(p_1, p_2) = (g_1(x_1, x_2, p_3), g_2(x_1, x_2, p_3))$ and Λ_s is the graph of a function $\nabla G(x_1, x_2, p_3)$ which means that $\frac{\partial G}{\partial x_1} = g_1$, $\frac{\partial G}{\partial x_2} = g_2$, $\frac{\partial G}{\partial p_3} = f$. Then $\frac{\partial g_1}{\partial p_3} = \frac{\partial f}{\partial x_1}$, $\frac{\partial g_2}{\partial p_3} = \frac{\partial f}{\partial x_2}$, $\frac{\partial f}{\partial x_2} = \frac{\partial g_2}{\partial x_1}$, $\frac{\partial g_1}{\partial x_2} = \frac{\partial g_2}{\partial x_1}$. In this new setting, $\pi(x_1, x_2, p_3) = (x_1, x_2, f(x_1, x_2, p_3))$ $\det d\pi = \frac{\partial f}{\partial p_3}$ and cusp caustics occur when

$$\frac{\partial f}{\partial p_3} = \frac{\partial^2 f}{\partial p_3^2} = 0, \quad \frac{\partial^3 f}{\partial p_3^3} \neq 0, \quad \text{and } \left\{ \nabla \frac{\partial f}{\partial p_3}, \nabla \frac{\partial^2 f}{\partial p_3^2} \right\} \text{ are linearly independent.} \quad (8)$$

Next, we parametrize the canonical relation C of F in terms of x_1, x_2 and p_3 ; $(\alpha_1, \alpha_2, \sqrt{1 - |\alpha|^2})$, the take off direction of the reflected ray, writing $\alpha = (\alpha_1, \alpha_2)$; and τ , the variable dual to time.

$$C = \left\{ (r_1(\cdot), r_2(\cdot), t_{inc}(\cdot) + t_{ref}(\cdot), \rho_1(\cdot), \rho_2(\cdot), \tau; x_1, x_2, f(\cdot), \right. \\ \left. -\tau(c_0^{-1}(\cdot)\alpha_1 + g_1(\cdot)), -\tau(c_0^{-1}(\cdot)\alpha_2 + g_2(\cdot)), -\tau(c_0^{-1}(\cdot)\sqrt{1 - |\alpha|^2} + p_3)) \right\},$$

where

$$\begin{aligned} f(\cdot) &= f(x_1, x_2, p_3); & r_j(\cdot) &= r_j(x_1, x_2, f(x_1, x_2, p_3), \alpha), j = 1, 2; \\ t_{inc}(\cdot) &= t_{inc}(x_1, x_2, p_3); & t_{ref}(\cdot) &= t_{ref}(x_1, x_2, f(x_1, x_2, p_3), \alpha); \\ \rho_j(\cdot) &= \rho_j(x_1, x_2, f(x_1, x_2, p_3), \alpha), j = 1, 2; & g_j(\cdot) &= g_j(x_1, x_2, p_3), j = 1, 2; \\ & & \text{and } c_0^{-1}(\cdot) &= c_0^{-1}(x_1, x_2, f(\cdot)). \end{aligned}$$

We have $\pi_R(x_1, x_2, p_3, \alpha_1, \alpha_2, \tau) = (x_1, x_2, f(\cdot), -\tau(c_0^{-1}(\cdot)\alpha_1 + g_1(\cdot)), -\tau(c_0^{-1}(\cdot)\alpha_2 + g_2(\cdot)), -\tau(c_0^{-1}(\cdot)\sqrt{1-|\alpha|^2} + p_3))$ and thus

$$d\pi_R = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \frac{\partial f}{\partial x_1} & \frac{\partial f}{\partial x_2} & \frac{\partial f}{\partial p_3} & 0 & 0 & 0 \\ A & B & C & -\tau c_0^{-1} & 0 & -c_0^{-1}(\cdot)\alpha_1 - g_1(\cdot) \\ D & E & F & 0 & -\tau c_0^{-1} & -c_0^{-1}(\cdot)\alpha_2 - g_2(\cdot) \\ G & H & I & \frac{\tau c_0^{-1}\alpha_1}{\sqrt{1-|\alpha|^2}} & \frac{\tau c_0^{-1}\alpha_2}{\sqrt{1-|\alpha|^2}} & -c_0^{-1}(\cdot)\sqrt{1-|\alpha|^2} - p_3 \end{bmatrix}$$

where $A = \frac{\partial(-\tau(c_0^{-1}(\cdot)\alpha_1 + g_1(\cdot)))}{\partial x_1}$; $B = \frac{\partial(-\tau(c_0^{-1}(\cdot)\alpha_1 + g_1(\cdot)))}{\partial x_2}$; $C = \frac{\partial(-\tau(c_0^{-1}(\cdot)\alpha_1 + g_1(\cdot)))}{\partial p_3}$; $D = \frac{\partial(-\tau(c_0^{-1}(\cdot)\alpha_2 + g_2(\cdot)))}{\partial x_1}$; $E = \frac{\partial(-\tau(c_0^{-1}(\cdot)\alpha_2 + g_2(\cdot)))}{\partial x_2}$; $F = \frac{\partial(-\tau(c_0^{-1}(\cdot)\alpha_2 + g_2(\cdot)))}{\partial p_3}$; $G = \frac{\partial(-\tau(c_0^{-1}(\cdot)\sqrt{1-|\alpha|^2} + p_3))}{\partial x_1}$; $H = \frac{\partial(-\tau(c_0^{-1}(\cdot)\sqrt{1-|\alpha|^2} + p_3))}{\partial x_2}$; and $I = \frac{\partial(-\tau(c_0^{-1}(\cdot)\sqrt{1-|\alpha|^2} + p_3))}{\partial p_3}$.

Hence $\det d\pi_R = \frac{\partial f}{\partial p_3}$ and the kernel at $\Sigma_1 = \{\det d\pi_R = 0\}$ is spanned by a vector v_1 which is $\frac{\partial f}{\partial p_3}$ plus a linear combination of $\{\frac{\partial}{\partial \alpha_1}, \frac{\partial}{\partial \alpha_2}, \frac{\partial}{\partial \tau}\}$, because the matrix

$$\begin{bmatrix} -\tau c_0^{-1} & 0 & c_0^{-1}(\cdot)\alpha_1 + g_1(\cdot) \\ 0 & -\tau c_0^{-1} & c_0^{-1}(\cdot)\alpha_2 + g_2(\cdot) \\ \frac{\tau c_0^{-1}\alpha_1}{\sqrt{1-|\alpha|^2}} & \frac{\tau c_0^{-1}\alpha_2}{\sqrt{1-|\alpha|^2}} & c_0^{-1}(\cdot)\sqrt{1-|\alpha|^2} + p_3 \end{bmatrix}$$

is nondegenerate by the calculations of [28]. From the cusp conditions (8), one can see that π_R has a cusp singularity.

Similarly, $\pi_L(x_1, x_2, \alpha_1, \alpha_2, p_3, \tau) = (r_1, r_2, t_{inc} + t_{ref}, \rho_1, \rho_2, \tau)$ and

$$d\pi_L = \begin{bmatrix} \frac{\partial r_1}{\partial x_1} + \frac{\partial r_1}{\partial x_3} \frac{\partial f}{\partial x_1} & \frac{\partial r_1}{\partial x_2} + \frac{\partial r_1}{\partial x_3} \frac{\partial f}{\partial x_2} & \frac{\partial r_1}{\partial \alpha_1} & \frac{\partial r_1}{\partial \alpha_2} & \frac{\partial r_1}{\partial p_3} & \frac{\partial f}{\partial x_3} \frac{\partial f}{\partial p_3} & 0 \\ \frac{\partial r_2}{\partial x_1} + \frac{\partial r_2}{\partial x_3} \frac{\partial f}{\partial x_1} & \frac{\partial r_2}{\partial x_2} + \frac{\partial r_2}{\partial x_3} \frac{\partial f}{\partial x_2} & \frac{\partial r_2}{\partial \alpha_1} & \frac{\partial r_2}{\partial \alpha_2} & \frac{\partial r_2}{\partial p_3} & \frac{\partial f}{\partial x_3} \frac{\partial f}{\partial p_3} & 0 \\ \frac{\partial t_{ref}}{\partial x_1} + \frac{\partial t_{inc}}{\partial x_1} & \frac{\partial t_{ref}}{\partial x_2} + \frac{\partial t_{inc}}{\partial x_2} & \frac{\partial t_{ref}}{\partial \alpha_1} & \frac{\partial t_{ref}}{\partial \alpha_2} & \frac{\partial t_{ref}}{\partial p_3} & \frac{\partial f}{\partial x_3} \frac{\partial f}{\partial p_3} + \frac{\partial t_{inc}}{\partial p_3} & 0 \\ \frac{\partial \rho_1}{\partial x_1} + \frac{\partial \rho_1}{\partial x_3} \frac{\partial f}{\partial x_1} & \frac{\partial \rho_1}{\partial x_2} + \frac{\partial \rho_1}{\partial x_3} \frac{\partial f}{\partial x_2} & \frac{\partial \rho_1}{\partial \alpha_1} & \frac{\partial \rho_1}{\partial \alpha_2} & \frac{\partial \rho_1}{\partial p_3} & \frac{\partial f}{\partial x_3} \frac{\partial f}{\partial p_3} & 0 \\ \frac{\partial \rho_2}{\partial x_1} + \frac{\partial \rho_2}{\partial x_3} \frac{\partial f}{\partial x_1} & \frac{\partial \rho_2}{\partial x_2} + \frac{\partial \rho_2}{\partial x_3} \frac{\partial f}{\partial x_2} & \frac{\partial \rho_2}{\partial \alpha_1} & \frac{\partial \rho_2}{\partial \alpha_2} & \frac{\partial \rho_2}{\partial p_3} & \frac{\partial f}{\partial x_3} \frac{\partial f}{\partial p_3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

From [28] we have that $\frac{\partial t_{inc}}{\partial p_3} = 0$, $\frac{\partial r_1}{\partial x_3} = 0$, $\frac{\partial r_2}{\partial x_3} = 0$, $\frac{\partial \rho_1}{\partial x_3} = 0$, $\frac{\partial \rho_2}{\partial x_3} = 0$, $\frac{\partial t_{ref}}{\partial x_3} <$

0, and that the matrix

$$\begin{bmatrix} \frac{\partial r_1}{\partial x_1} & \frac{\partial r_1}{\partial x_2} & \frac{\partial r_1}{\partial \alpha_1} & \frac{\partial r_1}{\partial \alpha_2} \\ \frac{\partial r_2}{\partial x_1} & \frac{\partial r_2}{\partial x_2} & \frac{\partial r_2}{\partial \alpha_1} & \frac{\partial r_2}{\partial \alpha_2} \\ \frac{\partial \rho_1}{\partial x_1} & \frac{\partial \rho_1}{\partial x_2} & \frac{\partial \rho_1}{\partial \alpha_1} & \frac{\partial \rho_1}{\partial \alpha_2} \\ \frac{\partial \rho_2}{\partial x_1} & \frac{\partial \rho_2}{\partial x_2} & \frac{\partial \rho_2}{\partial \alpha_1} & \frac{\partial \rho_2}{\partial \alpha_2} \end{bmatrix}$$

is nondegenerate. Thus, $\det d\pi_L = \frac{\partial f}{\partial p_3}$, $\ker \pi_L = \frac{\partial}{\partial p_3}$ and as before one sees that π_L has also a cusp singularity.

Now we will show that the images of the cusp points are involutive. By Remark 1 in Sec. 2.5, it is enough to examine the degenerate symplectic form $\omega_C = \pi_R^* \omega_{T^*Y}$ on C and its kernel, equal to $\{v : \omega_C(v, w) = 0, \forall w \in TC\}$. We have from (8) that $\Sigma_1 = \{(x, \xi); f_{p_3} = 0\}$ and $\Sigma_{1,1} = \{(x, \xi); f_{p_3} = f_{p_3 p_3} = 0\}$. Since $\omega_C|_{\Sigma_1}$ drops rank by 2 and $\ker d\pi_R = \mathbb{R} \cdot v_1$, one has $\ker \omega_C = \text{span}\{v_1, v_2\}$, where v_2 is a vector, without $\frac{\partial}{\partial p_3}$ component, that we now determine. At Σ_1 , one has

$$\begin{aligned} \omega_C|_{\Sigma_1} &= (c_0^{-1}\alpha_1 + g_1 + (c_0^{-1}\sqrt{1-|\alpha|^2} + p_3)f_{x_1})(d\tau \wedge dx_1) \\ &+ (c_0^{-1}\alpha_2 + g_2 + (c_0^{-1}\sqrt{1-|\alpha|^2} + p_3)f_{x_2})(d\tau \wedge dx_2) \\ &+ \tau c_0^{-1}(1 - \frac{\alpha_1}{\sqrt{1-|\alpha|^2}}f_{x_1})(d\alpha_1 \wedge dx_1) + \tau c_0^{-1}(1 - \frac{\alpha_2}{\sqrt{1-|\alpha|^2}}f_{x_2})(d\alpha_2 \wedge dx_2) \\ &+ \tau c_0^{-1} \frac{-\alpha_1}{\sqrt{1-|\alpha|^2}}f_{x_2}(d\alpha_1 \wedge dx_2) + \tau c_0^{-1} \frac{-\alpha_2}{\sqrt{1-|\alpha|^2}}f_{x_1}(d\alpha_2 \wedge dx_1) \\ &+ \tau(\partial_{x_2}c_0^{-1}\alpha_1 - \partial_{x_1}c_0^{-1}\alpha_2 + \partial_{x_2}c_0^{-1}\sqrt{1-|\alpha|^2}f_{x_1} - \partial_{x_1}c_0^{-1}\sqrt{1-|\alpha|^2}f_{x_2})(dx_2 \wedge dx_1) \\ &+ 2\tau f_{x_1}(dp_3 \wedge dx_1) + 2\tau f_{x_2}(dp_3 \wedge dx_2). \end{aligned}$$

We need to look at the 5×5 skew symmetric matrix of ω_C corresponding to $x_1, x_2, \alpha_1, \alpha_2, \tau$:

$$\begin{bmatrix} 0 & a & \star & \star & -b \\ -a & 0 & \star & \star & -c \\ \tau c_0^{-1}(1 - \frac{\alpha_1}{\sqrt{1-|\alpha|^2}}f_{x_1}) & \tau c_0^{-1} \frac{-\alpha_1}{\sqrt{1-|\alpha|^2}}f_{x_2} & 0 & 0 & 0 \\ \tau c_0^{-1} \frac{-\alpha_2}{\sqrt{1-|\alpha|^2}}f_{x_1} & \tau c_0^{-1}(1 - \frac{\alpha_2}{\sqrt{1-|\alpha|^2}}f_{x_2}) & 0 & 0 & 0 \\ b & c & 0 & 0 & 0 \end{bmatrix}$$

where a is the coefficient of $dx_1 \wedge dx_2$, b is the coefficient of $d\tau \wedge dx_1$ and c is the coefficient of $d\tau \wedge dx_2$. v is a combination of $\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \frac{\partial}{\partial \alpha_1}, \frac{\partial}{\partial \alpha_2}, \frac{\partial}{\partial \tau}$; note

that

$$\left| \begin{array}{cc} 1 - \frac{\alpha_1}{\sqrt{1-|\alpha|^2}} f_{x_1} & \frac{-\alpha_1}{\sqrt{1-|\alpha|^2}} f_{x_2} \\ \frac{-\alpha_2}{\sqrt{1-|\alpha|^2}} f_{x_1} & 1 - \frac{\alpha_2}{\sqrt{1-|\alpha|^2}} f_{x_2} \end{array} \right| = 1 + \nabla \sqrt{1-|\alpha|^2} \cdot \nabla_{x_1, x_2} f \neq 0,$$

which means that v_2 does not contain any $\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}$ terms, but only $\frac{\partial}{\partial \alpha_1}, \frac{\partial}{\partial \alpha_2}, \frac{\partial}{\partial \tau}$. This shows that $\ker \omega_C$ is simply tangent to Σ_1 at $\Sigma_{1,1}$, and thus $\pi_R(\Sigma_{1,1})$ is involutive; by Remark 1, so is $\pi_L(\Sigma_{1,1})$.

4 Weak normal forms for flat two-sided cusps

Next, we show that any flat two-sided cusp can be prepared, by application of suitable canonical transformations on the left and right, so as to be parametrized by a phase function similar to that for R_0 in (2). Recall that one can conjugate any two-sided *fold* to a single normal form [22, 23], but, as in [16, 7], we will need to work with merely approximate normal forms. For simplicity, we restrict ourselves to the three dimensional setting of interest for the seismic problem.

Assume $\dim(X) = \dim(Y) = 3$ and suppose $C \subset T^*X \times T^*Y$ is a flat two-sided cusp. Let $c^0 \in \Sigma_{1,1}$. We claim that there exist local canonical coordinates (x, ξ) on T^*X near $(x^0, \xi^0) := \pi_L(c^0)$ and (y, η) on T^*Y near $(y^0, \eta^0) := \pi_R(c^0)$ such that, as for the model C_0 in Sec. 2, $(x_1, x', y_1, \eta') := (x_1, x_2, x_3, y_1, \eta_2, \eta_3)$ form local coordinates on C near c^0 . In fact, since $d\pi_L$ and $d\pi_R$ drop rank by 1 at c^0 , there exist (see [19]) symplectic decompositions $T_{(x^0, \xi^0)}(T^*X) = V_1 \oplus V'$, $T_{(y^0, \eta^0)}(T^*Y) = W_1 \oplus W'$, with $\dim(V_1) = \dim(W_1) = 2$, $\dim(V') = \dim(W') = 4$, and

$$T_{c^0}C = Gr(\chi) \oplus (\Lambda_L \times (0)) \oplus ((0) \times \Lambda_R),$$

where $\Lambda_L \subset V_1$, $\Lambda_R \subset W_1$ are Lagrangian, i.e., one-dimensional, subspaces and $\chi \in Sp(V', W')$. Thus, in suitable linear symplectic coordinates, $\Lambda_L = \{\xi_1 = 0\}$, $\Lambda_R = \{\eta_1 = 0\}$ in $V_1 \simeq W_1 \simeq T^*\mathbb{R}$, and $\chi = I \in Sp(T^*\mathbb{R}^2, T^*\mathbb{R}^2)$, so that (x_1, y_1, x', η') are coordinates on $T_{c^0}C$. By a standard application of the homogeneous Darboux theorem [19, 22], there are local canonical coordinates (x, ξ, y, η) on $T^*X \times T^*Y$ such that $c^0 = (0, e_3^*; 0, e_3^*)$ and (x, y_1, η') form local coordinates on C .

We further prepare C by noting that if v_R, v_L are kernel vector fields for π_R, π_L , i.e., nonzero vector fields along Σ_1 generating $\ker(d\pi_R)$, $\ker(d\pi_L)$,

resp., then v_L is a linear combination of $\partial_{y_1}, \partial\eta_2, \partial\eta_3$. Thus, $d\pi_R(v_L)$ is a vector field along $\pi_R(\Sigma_{1,1})$, nontangent to $\Sigma_{1,1}$ since π_R is a cusp. Using homogeneous Darboux and the nonradiality of $\pi_R(\Sigma_{1,1})$, one can assume that $v_L = \partial_{y_1}$ at $\Sigma_{1,1}$, $\pi_R(\Sigma_{1,1}) = \{\eta_1 = \eta_2 = 0\}$ and $d\pi_R(TC) = \text{span}\{T\pi_R(\Sigma_{1,1}), d\pi_R(v_L)\} = \{d\eta_1 = 0\}$ along $\Sigma_{1,1}$. Working similarly on the left, one can assume $v_R = \partial_{x_1}$, $\pi_L(\Sigma_{1,1}) = \{\xi_1 = \xi_2 = 0\}$ and $d\pi_L(TC) = \{d\xi_1 = 0\}$ along $\Sigma_{1,1}$.

Furthermore, near c^0 one can assume that (i) $\Sigma_1 = \{\eta_2 = f(x, y_1, \eta_3)\}$, and (ii) $\Sigma_{1,1} = \{\eta_2 = f(x, y_1, \eta_3), y_1 = g(x, \eta_3)\}$ for functions f, g , homogeneous of degrees 1, 0, resp., and with $\partial_{x_1}g \neq 0$. To see (i), let $\tilde{f}(x, y_1, \eta')$ be a defining function for Σ_1 , homogeneous of degree 1; by the implicit function theorem it suffices to show that $d_{\eta_2}\tilde{f}(c^0) \neq 0$. Euler's identity implies that $\eta' \cdot d_{\eta'}\tilde{f} = 0$ at Σ_1 ; since $\eta_2 = 0$, $\eta_3 \neq 0$ at c^0 , this yields $d_{\eta_3}\tilde{f} = 0$. Since π_L, π_R are cusps, the kernel vector fields $v_L, v_R \in T\Sigma_1$ at $\Sigma_{1,1}$, thus at c^0 , and hence $d_{x_1}\tilde{f}(c^0) = d_{y_1}\tilde{f}(c^0) = 0$. If $d_{\eta_2}\tilde{f} = 0$, the only nonzero components of $d\tilde{f}(c^0)$ would be $d_{x'}\tilde{f}$, and this contradicts the fact that $d\pi_L(T\Sigma_1) = d\pi_L(T\Sigma_{1,1}) = T\pi_L(\Sigma_{1,1}) = \{d\xi_1 = d\xi_2 = 0\}$. Hence, $d_{\eta_2}\tilde{f}(c^0) \neq 0$, and thus Σ_1 can be described as $\{\eta_2 = f(x, y_1, \eta_3)\}$ for some f . Note that since $v_R, v_L \in T\Sigma_1$ at $\Sigma_{1,1}$, one has $\partial_{x_1}f|_{\Sigma_{1,1}} = \partial_{y_1}f|_{\Sigma_{1,1}} = 0$. For (ii), note that since $\pi_L : \Sigma_{1,1} \rightarrow \pi_L(\Sigma_{1,1})$ is a diffeomorphism, and (x, ξ_3) are coordinates on $\pi_L(\Sigma_{1,1})$, it follows that $(\pi_L^*(x), \pi_L^*(\xi_3)) = (x, d_{x_3}S(x, y_1, \eta'))$ form coordinates on $\Sigma_{1,1}$. Since $T\Sigma_1 = \text{span}\{T\Sigma_{1,1}, v\}$ for both $v = v_R = \partial_{x_1}$ and $v = v_L = \partial_{y_1}$, $(x, y_1, d_{x_3}S(x, y_1, f, \eta_3))$ are coordinates on Σ_1 , and we can replace $d_{x_3}S$ by η_3 since $d_{x_3\eta_3}^2 S \neq 0$. Then, $\Sigma_{1,1}$ is a hypersurface in Σ_1 transverse to both ∂_{x_1} and ∂_{y_1} , and can thus be described as $\{\eta_2 = f(x, y_1, \eta_3), y_1 = g(x, \eta_3)\}$ with $\partial_{x_1}g \neq 0$.

Since (x, y_1, η') form microlocal coordinates on C , there exists a generating function $S(x, y_1, \theta')$, so that $\phi(x, y, \theta') = S(x, y_1, \theta') - y' \cdot \theta'$ parametrizes C near c^0 . We have

$$C = \left\{ (x, \partial_x S; y_1, \partial_{\theta'} S, -\partial_{y_1} S, \theta') : x \in \mathbb{R}^3, y_1 \in \mathbb{R}, \theta' \in \mathbb{R}^2 \setminus 0 \right\},$$

with $\pi_L(x, y_1, \theta') = (x, \partial_x S(x, y_1, \theta'))$, $\pi_R(x, y_1, \theta') = (y, -\partial_{y_1} S(x, y_1, \theta'), \theta')$ and $\Sigma_{1,1} = \{\theta_2 - f = y_1 - g = 0\}$. Since $\pi_L(\Sigma_{1,1}) = \{\xi_1 = \xi_2 = 0\}$, one has $\partial_{x_1}S|_{\{\theta_2 - f = y_1 - g = 0\}} = 0$ and $\partial_{x_2}S|_{\{\theta_2 - f = y_1 - g = 0\}} = 0$. From the second relation, it follows that $S|_{\{\theta_2 - f = y_1 - g = 0\}}$ is independent of x_2 , so that S is a function of just x_1, x_3, θ_3 ; however, from the first relation one can express S

as

$$S(x, y_1, \theta) = S_0(x_3, \theta_3) + (y_1 - g)^2 S_1(x, y_1, \theta') + (\theta_2 - f) S_2(x, y_1, \theta')$$

Similarly, $\pi_R(\Sigma_{1,1}) = \{\eta_1 = \eta_2 = 0\}$ implies that $\partial_{y_1} S|_{\{\theta_2 - f = 0 = y_1 - g\}} = 0$, but doesn't provide any more information about S , cf. Remark 1. On C , the ideal of $\Sigma_{1,1}$ equals $(\theta_2 - f(x, y_1, \theta_3), y_1 - g(x, \theta_3)) = (S_{y_1}, \theta_2)$, so $f = \alpha S_{y_1} + \beta \theta_2$. Hence $f|_{\Sigma_{1,1}} = 0$; $\partial_{\theta_3} f|_{\Sigma_{1,1}} = 0$ (since f is homogeneous of degree 1); $f = c(y_1 - g)^2$ for some $c \in C^\infty$; and $\partial_{\theta_3} g|_{\Sigma_{1,1}} = 0$, since g is homogeneous of degree 0. Thus, the canonical relation must have the form,

$$\begin{aligned} C = \bigg\{ & (x_1, x_2, x_3, -2(y_1 - g)\partial_{x_1} g S_1 + (y_1 - g)^2 \partial_{x_1} S_1 + (\theta_2 - f)\partial_{x_1} S_2 - \partial_{x_1} f S_2, \\ & (y_1 - g)^2 \partial_{x_2} S_1 - 2(y_1 - g)\partial_{x_2} g S_1 + (\theta_2 - f)\partial_{x_2} S_2 - \partial_{x_2} f S_2, \\ & (y_1 - g)^2 \partial_{x_3} S_1 - 2(y_1 - g)\partial_{x_3} g S_1 + (\theta_2 - f)\partial_{x_3} S_2 - \partial_{x_3} f S_2 + \partial_{x_3} S_0; \\ & y_1, y_2, y_3, 2(y_1 - g)S_1 + (y_1 - g)^2 \partial_{y_1} S_1 + (\theta_2 - f)\partial_{y_1} S_2 - \partial_{y_1} f S_2, \\ & -\theta_2, -\theta_3) : y_2 = (y_1 - g)^2 \partial_{\theta_2} S_1 + (\theta_2 - f)\partial_{\theta_2} S_2 + S_2, \\ & y_3 = (y_1 - g)^2 \partial_{\theta_3} S_1 - 2(y_1 - g)\partial_{\theta_3} g S_1 + (\theta_2 - f)\partial_{\theta_3} S_2 - \partial_{\theta_3} f S_2 + \partial_{\theta_3} S_3 \bigg\}. \end{aligned}$$

From this, one sees that

$$d\pi_L \equiv \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ \cdot & \cdot & \cdot & -2\partial_{x_1} g S_1 - \partial_{x_1 y_1} f S_2 & \partial_{x_1} S_2 & 0 \\ \cdot & \cdot & \cdot & -2\partial_{x_2} g S_1 - \partial_{x_2 y_1} f S_2 & \partial_{x_2} S_2 & 0 \\ \cdot & \cdot & \cdot & \cdot & \partial_{x_3} S_2 & \partial_{x_3 \theta_3}^2 S_0 \end{bmatrix} \text{ mod } (y_1 - g, \theta_2 - f)$$

with $\det d\pi_L = [(-2\partial_{x_1} g S_1 - \partial_{x_1 y_1} f S_2)\partial_{x_2} S_2 - (-2\partial_{x_2} g S_1 - \partial_{x_2 y_1} f S_2)\partial_{x_1} S_2](\partial_{x_3 \theta_3}^2 S_0) + (y_1 - g)E_1 + (\theta_2 - f)E_2$ for some E_1, E_2 . Since π_L has a cusp singularity at $\Sigma_{1,1}$, the determinant must vanish simply. We have that $\partial_{x_3 \theta_3}^2 S_0|_{\Sigma_{1,1}} \neq 0$ by homogeneity and $\partial_{x_2 \theta_2} S = \partial_{x_2} S_2|_{\Sigma_{1,1}} \neq 0$ since $d\pi_L(TC) = \{d\xi_1 = 0\}$. Since $\text{Ker } d\pi_L = \partial_{y_1}$, the y_1 column in $d\pi_L$ must be 0 at $\Sigma_{1,1}$. We have that $(-2\partial_{x_1} g S_1 - \partial_{x_1 y_1} f S_2)|_{\Sigma_{1,1}} = \partial_{x_1} g(S_1 + cS_2)|_{\Sigma_{1,1}} = 0$ so $(S_1 + cS_2)|_{\Sigma_{1,1}} = 0$, and $(-2\partial_{x_2} g S_1 - \partial_{x_2 y_1} f S_2)|_{\Sigma_{1,1}} = \partial_{x_2} g(S_1 + cS_2)|_{\Sigma_{1,1}} = 0$, and since $\partial_{x_2} S_2, \partial_{x_1} g \neq 0$ we can conclude that $\det d\pi_L|_{\Sigma_{1,1}} = (S_1 + cS_2)|_{\Sigma_{1,1}}$. Since S_0 is nondegenerate, by another canonical transformation on the left we may assume that $S_0 = x_3 \theta_3$. From $\ker d\pi_L = \mathbb{R} \cdot \frac{\partial}{\partial y_1}$, we have $\partial_{y_1}(S_1 + cS_2)|_{\Sigma_{1,1}} = 0$. Since $d\pi_R(TC) =$

$\{d\eta_1 = 0\}$ we obtain that $\partial_{y_1} S_2|_{\Sigma_{1,1}} = \partial_{\theta_2 y_1} S|_{\Sigma_{1,1}} = 0$. One also has that $\partial_{y_1} S_1|_{\Sigma_{1,1}} = 0$, since a calculation shows that this is $\partial_{y_1}^2 \eta_1(x, y_1, \theta')|_{\Sigma_{1,1}}$, which vanishes due to the cusp structure of $d\pi_R$ and the fact that $d\pi_R(TC) = \{d\eta_1 = 0\}$. Then one can write $S_1 = (y_1 - g)^2 S_5 + (\theta_2 - f) S_6$ with $S_5 \neq 0$, leading to a weak normal form for a phase function parametrizing the flat two-sided cusp C , namely $\phi(x, y, \theta_2, \theta_3) = (S_2 + (y_1 - g)^2 S_6)(\theta_2 - f) + (x_3 - y_3)\theta_3 + (y_1 - g)^4 S_5 - y_2 \theta_2$, with $\partial_{x_2} S_2(x, y_1, \theta')|_{\Sigma_{1,1}} \neq 0$, $S_5, S_6 \neq 0$.

In studying the composition in the next section, f, g introduce only algebraic complications and do not affect the final result. Thus, for simplicity we take $f = 0$ and $g = x_1$ and will use the phase function

$$\begin{aligned} \phi(x, y, \theta_2, \theta_3) &= (S_2 - y_2 + (x_1 - y_1)^2 S_4) \theta_2 + (x_3 - y_3) \theta_3 + (x_1 - y_1)^4 S_3, \\ \partial_{x_2} S_2(x, y_1, \theta') &\neq 0, \quad S_3, S_4 \neq 0 \text{ on } \Sigma_{1,1}, \end{aligned} \quad (9)$$

with S_5, S_6 relabeled as S_3, S_4 , resp.

5 Composition

Now consider the composition B^*A for $A \in I^m(X, Y; C)$, $B \in I^{m'}(X, Y; C)$. Conjugating by unitary FIOs associated with the canonical transformations used in the last section, we may assume that C is parametrized by the phase function ϕ from (9). We have, for $a \in S^{m+\frac{1}{2}}$, $b \in S^{m'+\frac{1}{2}}$,

$$B^*A f(x) = \int e^{i(\phi(z, y, \eta') - \phi(z, x, \theta'))} a(z, y, \eta') \bar{b}(z, x, \theta') f(y) dy dz d\theta' d\eta',$$

where the phase $\phi(z, y, \eta') - \phi(z, x, \theta')$, which we denote $\widehat{\phi}$, is

$$\begin{aligned} (z_3 - y_3)\eta_3 &+ (z_1 - y_1)^4 S_3(z, y_1, \eta') + (S_2(z, y_1, \eta') - y_2 + (z_1 - y_1)^2 S_4(z, y_1, \eta')) \eta_2 \\ - (z_3 - x_3)\theta_3 &- (z_1 - x_1)^4 S_3(z, x_1, \theta') - (S_2(z, x_1, \theta') - x_2 + (z_1 - x_1)^2 S_4(z, x_1, \theta')) \theta_2. \end{aligned}$$

We perform stationary phase in z_2, η_2 and z_3, η_3 :

$$\begin{aligned} \partial_{z_2} \widehat{\phi} &= \eta_2 \partial_{z_2} S_2 - \theta_2 \partial_{z_2} S_2 + \partial_{z_2} S_3 (z_1 - y_1)^4 - \partial_{z_2} S_3 (z_1 - x_1)^4 \\ &+ (z_1 - y_1)^2 \eta_2 \partial_{z_2} S_4 - (z_1 - x_1)^2 \theta_2 \partial_{z_2} S_4, \end{aligned} \quad (10)$$

$$\begin{aligned} \partial_{z_3} \widehat{\phi} &= \eta_3 - \theta_3 + (z_1 - y_1)^4 \partial_{z_3} S_3 - (z_1 - x_1)^4 \partial_{z_3} S_3 + \eta_2 \partial_{z_3} S_2 - \theta_2 \partial_{z_3} S_2 \\ &+ (z_1 - y_1)^2 \eta_2 \partial_{z_3} S_4 - (z_1 - x_1)^2 \theta_2 \partial_{z_3} S_4, \end{aligned} \quad (11)$$

$$\partial_{\eta_2} \widehat{\phi} = (z_1 - y_1)^4 \partial_{\eta_2} S_3 + S_2 - y_2 + (z_1 - y_1)^2 [S_4 + \eta_2 \partial_{\eta_2} S_4] + \eta_2 \partial_{\eta_2} S_2 \quad (12)$$

$$\partial_{\eta_3} \widehat{\phi} = z_3 - y_3 + (z_1 - y_1)^4 \partial_{\eta_3} S_3 + \eta_2 \partial_{\eta_3} S_2 + (z_1 - y_1)^2 \eta_2 \partial_{\eta_3} S_4. \quad (13)$$

We solve for z_3 using (13) and find

$$z_3 = y_3 - (z_1 - y_1)^4 \partial_{\eta_3} S_3 - \eta_2 \partial_{\eta_3} S_2 - (z_1 - y_1)^2 \eta_2 \partial_{\eta_3} S_4 := y_3 + S_5.$$

From equation (12) we can solve for z_2 since $\partial_{z_2} S_2 \neq 0$, but only implicitly. Now rewrite (11) as

$$\begin{aligned} \eta_3 = & \theta_3 + (z_1 - x_1)^4 \partial_{z_3} S_3 - (z_1 - y_1)^4 \partial_{z_3} S_3 \\ & + (z_1 - x_1)^2 \theta_2 \partial_{z_3} S_4 - (z_1 - y_1)^2 \eta_2 \partial_{z_3} S_4 + \theta_2 \partial_{z_3} S_2 - \eta_2 \partial_{z_3} S_2, \end{aligned}$$

and solve for η_3 , using the fact that $\theta_2 \partial_{z_3} S_2 - \eta_2 \partial_{z_3} S_2$ vanishes at $\{x_1 = y_1, \eta' = \theta'\}$, so that

$$\theta_2 \partial_{z_3} S_2 - \eta_2 \partial_{z_3} S_2 = (x_1 - y_1) \theta_2 \partial_{z_3 x_1}^2 S_2 + (\theta_2 - \eta_2) (\partial_{z_3} S_2 + \theta_2 \partial_{z_3 \theta_2}^2 S_2) + (\theta_3 - \eta_3) \theta_2 \partial_{z_3 \theta_3}^2 S_2.$$

The other differences in the equations (10),(13) can be handled similarly, so

$$\begin{aligned} \eta_3 = & \theta_3 + (x_1 - y_1) M_1(z_1, z_2, x_1, y_1, \theta', \eta_2) + (\theta_2 - \eta_2) N_1(z_1, z_2, x_1, y_1, \theta', \eta_2) + \\ & (\theta_3 - \eta_3) P(z_1, z_2, x_1, y_1, \theta', \eta_2) \end{aligned}$$

for some functions M_1, N_1 and P . Solving for $\eta_3 - \theta_3$, we have

$$\eta_3 - \theta_3 = [(x_1 - y_1) M_1 + (\theta_2 - \eta_2) N_1] (I + P)^{-1} = (x_1 - y_1) M_2 + (\theta_2 - \eta_2) N_2.$$

Also, $\eta_2 - \theta_2 = (x_1 - y_1) M_3 + (\eta_3 - \theta_3) N_3$, so that $\eta_3 - \theta_3 = (x_1 - y_1) M_4$ and $\eta_2 - \theta_2 = (x_1 - y_1) N_4$. Hence, the phase becomes

$$\begin{aligned} \widehat{\phi} = & (x_3 - y_3) \theta_3 + (x_2 - y_2) \theta_2 + ((z_1 - y_1)^2 S_4(z_1, y, \theta') - (z_1 - x_1)^2 S_4(z_1, x_1, y_2, y_3, \theta')) \theta_2 \\ & + (z_1 - y_1)^4 S_3 - (z_1 - x_1)^4 S_3 + (S_2(z_1, y, \theta') - S_2(z_1, x_1, y_2, y_3, \theta')) \theta_2 + (x_1 - y_1) S_6. \end{aligned}$$

We have

$$\begin{aligned} ((z_1 - y_1)^2 S_4(z_1, y, \theta') - (z_1 - x_1)^2 S_4(z_1, x_1, y_2, y_3, \theta')) \theta_2 &= [(z_1 - y_1)^2 S_4(z_1, y_1, \cdot) - (z_1 - x_1)^2 S_4(z_1, y_1, \cdot)] \theta_2 \\ &+ [(z_1 - x_1)^2 S_4(z_1, y_1, \cdot) - (z_1 - x_1)^2 S_4(z_1, x_1, \cdot)] \theta_2 \\ &= [(x_1 - y_1)(2z_1 - x_1 - y_1) S_4(z_1, y_1, \cdot) + (z_1 - x_1)^2 (x_1 - y_1) \partial_{y_1} S_4] \theta_2, \end{aligned}$$

$$\begin{aligned}
(z_1 - y_1)^4 S_3 - (z_1 - x_1)^4 S_3 &= (z_1 - y_1)^4 S_3(z_1, y_1, \cdot) - (z_1 - x_1)^4 S_3(z_1, y_1, \cdot) \\
&\quad + (z_1 - x_1)^4 S_3(z_1, y_1, \cdot) - (z_1 - x_1)^4 S_3(z_1, x_1, \cdot) \\
&= (x_1 - y_1)(2z_1 - x_1 - y_1)((z_1 - y_1)^2 + (z_1 - x_1)^2) S_3(z_1, y_1, \cdot) \\
&\quad + (z_1 - x_1)^4 (x_1 - y_1) \partial_{y_1} S_3,
\end{aligned}$$

and can write $(z_1 - x_1)^2 (x_1 - y_1) \partial_{y_1} S_4 \theta_2 + (z_1 - x_1)^4 (x_1 - y_1) \partial_{y_1} S_3 + (S_2(z_1, y, \theta') - S_2(z_1, x_1, y_2, y_3, \theta')) \theta_2 + (x_1 - y_1) S_6 := (x_1 - y_1) S_7$. From this, it follows that $\widehat{\phi} = (x_2 - y_2) \theta_2 + (x_3 - y_3) \theta_3 + (x_1 - y_1) [(2z_1 - x_1 - y_1)((z_1 - y_1)^2 + (z_1 - x_1)^2) S_3 + (2z_1 - x_1 - y_1) S_4 \theta_2 + S_7(z_1, y, x_1, \theta)]$. We make a change of variables similar to that in the model case in Sec. 2.4:

$$\theta_1 = (2z_1 - x_1 - y_1)((z_1 - y_1)^2 + (z_1 - x_1)^2) S_3 + (2z_1 - x_1 - y_1) S_4 \theta_2 + S_7,$$

$$\begin{aligned}
\frac{\partial \theta_1}{\partial z_1} &= 2S_4 \theta_2 + (2z_1 - x_1 - y_1) \theta_2 \partial_{z_1} S_4 + 2S_3((z_1 - x_1)^2 + (z_1 - y_1)^2) \\
&\quad + 2(2z_1 - x_1 - y_1)^2 S_3 + (2z_1 - x_1 - y_1)((z_1 - y_1)^2 + (z_1 - x_1)^2) \partial_{z_1} S_3 + \partial_{z_1} S_7
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial^2 \theta_1}{\partial z_1^2} &= 4\partial_{z_1} S_4 \theta_2 + (2z_1 - x_1 - y_1) \theta_2 \partial_{z_1}^2 S_4 + 4((z_1 - x_1)^2 + (z_1 - y_1)^2) \partial_{z_1} S_3 \\
&\quad + 12(2z_1 - x_1 - y_1) S_3 + 4(2z_1 - x_1 - y_1)^2 \partial_{z_1} S_3 + (2z_1 - x_1 - y_1)((z_1 - y_1)^2 \\
&\quad + (z_1 - x_1)^2) \partial_{z_1}^2 S_3 + \partial_{z_1} S_7.
\end{aligned}$$

Near points on $\Sigma_{1,1}$, $\frac{\partial^2 \theta_1}{\partial z_1^2} = 0$ implies $z_1 = \frac{x_1 + y_1}{2} + R(x_1, y, \theta_2, \theta_3)$, for some R which vanishes at the base point $(x_1, y, \theta_2, \theta_3) = (0, 0, 0, 0, 0, 1)$. Then

$$\begin{aligned}
\frac{\partial \theta_1}{\partial z_1} &= 2S_4 \theta_2 + R \partial_{z_1} S_4 \theta_2 + 2S_3 \left(\frac{(x_1 - y_1)^2}{2} + 2R^2 \right) \\
&\quad + 2R^2 S_3 + R \left(\frac{(x_1 - y_1)^2}{2} + 2R^2 \right) \partial_{z_1} S_3 + \partial_{z_1} S_7 = 0
\end{aligned}$$

implies that

$$\theta_2 = -\frac{(x_1 - y_1)^2}{2} \frac{2S_3 + R \partial_{z_1} S_3}{2S_4 + R \partial_{z_1} S_4} + P_1(x_1, y, \theta_3) = -\frac{(x_1 - y_1)^2}{2} N(x_1, y, \theta_3) + P_1(x_1, y, \theta_3)$$

for some N, P_1 , and

$$\theta_1 = R\left(\frac{(x_1 + y_1)^2}{2} + 2R^2\right)S_3 + RS_4\theta_2 + S_7 := P_2(x_1, y, \theta_3).$$

Since $S_3, S_4 \neq 0$ and $R = 0$ at the base point, one has $N \neq 0$. Following the same analysis as in the model case, one sees that

$$\begin{aligned} \widehat{\phi} &= (x_2 - y_2)\theta_2 + (x_3 - y_3)\theta_3 + (x_1 - y_1)\theta_1 + \left(\frac{\theta_1 - P_2}{\theta_3} - t_1t_2 - \frac{1}{2}t_1^3\right)\sigma_1 + \\ &\quad \left(\frac{\theta_2}{\theta_3} + \frac{(x_1 - y_1)^2}{2}\frac{N}{\theta_3} - \frac{P_1}{\theta_3} - t_2\right)\sigma_2. \end{aligned}$$

Next, we perform a stationary phase in t_2, σ_2 :

$$\partial_{t_2}\widehat{\phi} = -t_1\sigma_1 - \sigma_2, \quad \partial_{\sigma_2}\widehat{\phi} = \frac{\theta_2}{\theta_3} + \frac{N}{2\theta_3}(x_1 - y_1)^2 - \frac{P_1}{\theta_3} - t_2$$

and

$$\widehat{\phi} = (x - y)\theta + \left(\frac{\theta_1}{\theta_3} - \frac{P_2}{\theta_3} - t_1\left[\frac{\theta_2}{\theta_3} + \frac{N}{2\theta_3}(x_1 - y_1)^2 - \frac{P_1}{\theta_3}\right] - \frac{1}{2}t_1^3\right)\sigma_1$$

With one more stationary phase in σ_1 and θ_1 , obtaining

$$\partial_{\theta_1}\widehat{\phi} = x_1 - y_1 + \frac{\sigma_1}{\theta_3}, \quad \partial_{\sigma_1}\widehat{\phi} = \frac{\theta_1}{\theta_3} - \frac{P_2}{\theta_3} - t_1\left[\frac{\theta_2}{\theta_3} + \frac{N}{2\theta_3}(x_1 - y_1)^2 - \frac{P_1}{\theta_3}\right] - \frac{1}{2}t_1^3,$$

and then homogenize t_1 by setting $\tau = t_1\theta_3$. The resulting phase is then

$$\begin{aligned} \widehat{\phi} &= \left(x_2 - y_2 + \frac{\tau}{\theta_3}(x_1 - y_1)\right)\theta_2 + \left(x_3 - y_3 + (x_1 - y_1)\frac{P_2 - \frac{\tau}{\theta_3}P_1}{\theta_3}\right) \\ &\quad + (x_1 - y_1)^3\frac{\tau}{\theta_3^2}\frac{N}{2} + (x_1 - y_1)\frac{1}{2}\left(\frac{\tau}{\theta_3}\right)^3\theta_3, \end{aligned} \quad (14)$$

and the Schwartz kernel has the representation

$$K_{F^*F}(x, y) = \int e^{i\widehat{\phi}} c(x, y, \theta', \tau) d\theta_2 d\theta_3 d\tau, \quad c \in S^{m+m'}. \quad (15)$$

For each of the variables $s = x_1, y_1, y_2, y_3$, let $[[s]]$ denote

$$[[s]] = \partial_s \left(P_2 - \frac{\tau}{\theta_3}P_1\right) + (x_1 - y_1)^2 \frac{\tau}{\theta_3} \partial_s \frac{N}{2}, \quad [[\theta_3]] = \partial_{\theta_3} \left(P_2 - \frac{\tau}{\theta_3}P_1 + (x_1 - y_1)^2 \frac{\tau}{\theta_3} \frac{N}{2} + \frac{1}{2} \frac{\tau^3}{\theta_3^2}\right).$$

Then we have

$$\begin{aligned}
WF(F^*F) \subset & \left\{ (x_1, x_2, x_3, \frac{\tau\theta_2}{\theta_3} + 3(x_1 - y_1)^2\tau\frac{N}{2\theta_3} + P_2 - \frac{\tau}{\theta_3}P_1 + \frac{1}{2}(\frac{\tau}{\theta_3})^3\theta_3 + (x_1 - y_1)[[x_1]], \theta'; \right. \\
& y_1, y_2, y_3, \frac{\tau\theta_2}{\theta_3} + 3(x_1 - y_1)^2\tau\frac{N}{2\theta_3} + P_2 - \frac{\tau}{\theta_3}P_1 + \frac{1}{2}(\frac{\tau}{\theta_3})^3\theta_3 + (x_1 - y_1)[[y_1]], \\
& \left. \theta_2 + (x_1 - y_1)[[y_2]], \theta_3 + (x_1 - y_1)[[y_3]]) : \right. \\
& d_\tau \hat{\phi} = (x_1 - y_1) \left(\frac{\theta_2 - P_1}{\theta_3} + \frac{3}{2}(\frac{\tau}{\theta_3})^2 + (x_1 - y_1)^2\frac{N}{2\theta_3} \right) = 0, \\
& d_{\theta_2} \hat{\phi} = x_2 - y_2 + (x_1 - y_1)\frac{\tau}{\theta_3} = 0, \\
& \left. d_{\theta_3} \hat{\phi} = x_3 - y_3 - (x_1 - y_1)\frac{\tau\theta_2}{\theta_3^2} + (x_1 - y_1)[[\theta_3]] = 0 \right\}.
\end{aligned}$$

The phase function is degenerate because the equation $d_\tau \hat{\phi} = 0$ has a normal crossing; from $\{x_1 - y_1 = 0\}$, the contribution to $WF(B^*A)$ is contained in

$$\begin{aligned}
\Delta = & \left\{ (x, \frac{\tau\theta_2}{\theta_3} + P_2 - \frac{\tau}{\theta_3}P_1 + \frac{1}{2}(\frac{\tau}{\theta_3})^3\theta_3, \theta_2, \theta_3; \right. \\
& \left. x, \frac{\tau\theta_2}{\theta_3} + P_2 - \frac{\tau}{\theta_3}P_1 + \frac{1}{2}(\frac{\tau}{\theta_3})^3\theta_3, \theta_2, \theta_3) : x \in \mathbb{R}^3, (\tau, \theta') \in \mathbb{R}^3 \setminus 0 \right\},
\end{aligned}$$

which is a cusp parametrization of Δ (as in the model case). On the other hand, if $x_1 - y_1 \neq 0$ then

$$\begin{aligned}
\theta_2 &= P_1 - \frac{3}{2}(\frac{\tau}{\theta_3})^2\theta_3 - (x_1 - y_1)^2\frac{N}{2}, \quad y_2 = x_2 + (x_1 - y_1)\frac{\tau}{\theta_3}, \\
y_3 &= x_3 - (x_1 - y_1)\frac{\tau\theta_2}{\theta_3^2} + (x_1 - y_1)[[\theta_3]], \\
\xi_1 &= P_2 - (\frac{\tau}{\theta_3})^3\theta_3 + \tau N \frac{(x_1 - y_1)^2}{\theta_3} + (x_1 - y_1)[[x_1]], \\
\eta_1 &= P_2 - (\frac{\tau}{\theta_3})^3\theta_3 + \tau N \frac{(x_1 - y_1)^2}{\theta_3} + (x_1 - y_1)[[y_1]], \\
\eta_2 &= \theta_2 + (x_1 - y_1)[[y_2]], \text{ and } \eta_3 = \theta_3 + (x_1 - y_1)[[y_3]].
\end{aligned}$$

Let $\tilde{C} := \tilde{C}_\phi$ be the image of $\psi(x_1, x_2, x_3, y_1, \theta_3, \tau) = (x, \xi; y, \eta)$ with $\xi_2 = \theta_2, \xi_3 = \theta_3$, and $\xi_1, \eta_1, \eta_2, \eta_3, y_2, y_3$ given above. As in the model case, ψ satisfies Def. 2.2: it drops rank by one simply at $\Sigma_1 = \{x_1 - y_1 = 0 = \tau\}$, and $\ker d\pi = \mathbb{R} \cdot \frac{\partial}{\partial \tau} \not\subseteq T\Sigma$, so that $\tilde{C}$ is an open umbrella.

5.1 Generalized Fourier integral operators for $\Delta \cup \tilde{C}$

Using the phase $\hat{\phi}$, one can now define a class of generalized FIOs, G , with the $WF(G)$ contained in the union of the diagonal and the open umbrella. For a general $\tilde{a} \in S^\mu$, define G by the right hand side of (15), but with the amplitude $\tilde{a}$; then one has $WF(G) \subset \Delta \cup \tilde{C}$. Furthermore, one can compute the orders of G on $\Delta \setminus \tilde{C}$ and $\tilde{C} \setminus \Delta$, as in the model case at the end of Sec. 2.4. Due to the normal crossing of $d_\tau \hat{\phi}$, the critical set $Crit_{\hat{\phi}} = \{(x, y, \tau, \theta') : d_\tau \hat{\phi} = 0, d_{\theta'} \hat{\phi} = 0\}$ decomposes as $Crit_\Delta \cup Crit_{\tilde{C}}$, and $\sigma = \rho(E_{\hat{\phi}})^{\frac{1}{2}}$, where $E_{\hat{\phi}} = \left| \frac{D(\lambda_i, \frac{\partial \hat{\phi}}{\partial \theta})}{D(x, \theta)} \right|^{-1}$ and λ_i are local coordinates on $Crit_{\hat{\phi}}$.

On $Crit_\Delta$, local coordinates are (x, τ, θ') and

$$E_{\hat{\phi}} = \left| \frac{D(x, \tau, \theta', \frac{\partial \hat{\phi}}{\partial \tau}, \frac{\partial \hat{\phi}}{\partial \theta_2}, \frac{\partial \hat{\phi}}{\partial \theta_3})}{D(x, \tau, \theta', y_1, y_2, y_3)} \right|^{-1} = \left(\frac{\theta_2}{\theta_3} + \frac{3}{2} \frac{\tau^2}{\theta_3^2} - \frac{P_1}{\theta_3} \right)^{-1};$$

hence $G \in I^\mu(\Delta \setminus \tilde{C})$ and the principal symbol satisfies $\sigma \sim \delta^{-\frac{1}{2}}$, where δ is the distance to $\Delta \cap \tilde{C}$. On the other hand, (x, y_1, τ, θ_3) are local coordinates on $Crit_{\tilde{C}}$,

$$\begin{aligned} E_{\hat{\phi}} &= \left| \frac{D(x, y_1, \tau, \theta_3, \frac{\partial \hat{\phi}}{\partial \tau}, \frac{\partial \hat{\phi}}{\partial \theta_2}, \frac{\partial \hat{\phi}}{\partial \theta_3})}{D(x, y_1, \tau, \theta_2, \theta_3, y_2, y_3, \theta_2)} \right|^{-1} \\ &= \left(\frac{x_1 - y_1}{\theta_3} + \frac{(x_1 - y_1)^2}{\theta_3} [\partial_{y_3 \theta_3} P_2 - \frac{\tau}{\theta_3} \partial_{y_3 \theta_3} P_1 + (x_1 - y_1)^2 \frac{\tau}{2\theta_3} \partial_{y_3 \theta_3} N] \right)^{-1} \\ &\sim \left(\frac{x_1 - y_1}{\theta_3} \right)^{-1}, \end{aligned}$$

$G \in I^\mu(\tilde{C} \setminus \Delta)$ and again $\sigma \sim \delta^{-\frac{1}{2}}$. The equality of the orders on Δ and $\tilde{C}$ away from $\Delta \cap \tilde{C}$ is consistent with the composition result in [28, 6] for two-sided folds. This gives a precise description of the nonremovable artifact in the linearized seismic inversion problem for the single source geometry in the presence of cusp caustics.

Acknowledgements

We would like to thank Cliff Nolan for patiently explaining some of the calculations in [28]. The second author was supported by NSF grants DMS-0551894 and -0853892.

References

- [1] T. Banchoff, T. Gaffney and C. McCrory, *Cusps of Gauss mappings*, Research Notes in Math., **55**, Pitman, Boston, 1982.
- [2] G. Beylkin, Imaging of discontinuities in the inverse problem by inversion of a generalized Radon transform, *Jour. Math. Phys.* **28** (1985), 99–108.
- [3] A. Comech and S. Cuccagna, Integral operators with two-sided cusp singularities, *Int. Math. Res. Not.* (2000), 1225–1242.
- [4] M. Christ, A. Nagel, E. Stein and S. Wainger, Singular and maximal Radon transforms: analysis and geometry, *Ann. of Math.* (2) **150** (1999), 489–577.
- [5] J.J. Duistermaat and V. Guillemin, The spectrum of positive elliptic operators and periodic bicharacteristics, *Invent. Math.* **29** (1975), 39–79.
- [6] R. Felea, Composition calculus of Fourier integral operators with fold and blowdown singularities, *Comm. P.D.E.* **30** (2005), 1717–1740.
- [7] R. Felea and A. Greenleaf, An FIO calculus for marine seismic imaging: folds and crosscaps, *Comm. P.D.E.* **33** (2008), 45–77.
- [8] R. Felea, A. Greenleaf and M. Pramanik, An FIO calculus for marine seismic imaging, II: Sobolev estimates, <http://arxiv.org/abs/0903.1780>; *Math. Annalen*, to appear.
- [9] A. Givental, Lagrangian imbeddings of surfaces and unfolded Whitney umbrella. (English) *Func. Anal. Appl.* **20**(3) (1986), 197–203
- [10] M. Golubitsky and V. Guillemin, *Stable mappings and their singularities*, Springer-Verlag, New York, 1973.
- [11] A. Greenleaf and A. Seeger, Fourier integral operators with fold singularities, *J. Reine Angew. Math.* **455** (1994), 35–56.
- [12] ———, ———, Fourier integral operators with cusp singularities, *Amer. Jour. Math.* **120** (1998), 1077–1119.

- [13] ———, ———, Oscillatory integral operators with low-order degeneracies, *Duke Math. Jour.* **112** (2002), 397–420.
- [14] ———, ———, Oscillatory and Fourier integral operators with degenerate canonical relations, in *Proc. 6th Int. Conf. on Harmonic Analysis and Partial Diff. Eqns. (El Escorial, 2000)*, Publ. Mat. 2002, Vol. Extra, 93–141.
- [15] A. Greenleaf and G. Uhlmann, Estimates for singular Radon transforms and pseudodifferential operators with singular symbols, *Jour. Func. Anal.* **89** (1990), 220–232.
- [16] ———, ———, Compositions of some singular Fourier integral operators and estimates for restricted X-ray transforms, *Ann. Inst. Fourier (Grenoble)* **40**(2) (1990), 443–466; II, *Duke Math. Jour.* **64** (1991), 415–444.
- [17] J. Guckenheimer, Caustics and non-degenerate Hamiltonians, *Topology* **13** (1974), 127–133.
- [18] V. Guillemin and G. Uhlmann, Oscillatory integrals with singular symbols, *Duke Math. Jour.* **48** (1981), 251–267.
- [19] L. Hörmander, *The Analysis of Linear Partial Differential Operators, III*, Grundlehren math. Wiss. **274**, Springer Verlag, Berlin, 1985.
- [20] A. ten Kroode, D. Smit and A. Verdel, A microlocal analysis of migration, *Wave Motion* **28** (1998), 149–172.
- [21] F. Marhuenda, Microlocal analysis of some isospectral deformations, *Trans. Amer. Math. Soc.* **343** (1994), 245–275.
- [22] R. Melrose, Equivalence of glancing hypersurfaces. II, *Math. Annalen* **255** (1981), 159–198.
- [23] R. Melrose and M. Taylor, *Near peak scattering and the corrected Kirchhoff approximation for a convex obstacle*, Adv. in Math. **55**(3) (1985), 242–315.
- [24] R. Melrose and G. Uhlmann, Lagrangian intersection and the Cauchy problem, *Comm. Pure Appl. Math.* **32**(4) (1979), 483–519.

- [25] ———, ———, Microlocal structure of involutive conical refraction, *Duke Math. J.* **46** (1979), 571–582.
- [26] G. Mendoza, Symbol calculus associated with intersecting Lagrangians, *Comm. P.D.E.* **7** (1982), 1035–1116.
- [27] B. Morin, Formes canoniques des singularites d’une application differentiable, *C.R. Acad. Sc. Paris*, **260** (1965) 5662–5665 and 6503–6506.
- [28] C. Nolan, Scattering in the presence of fold caustics, *SIAM J. Appl. Math.* **61** (2000), 659–672.
- [29] C. Nolan and W. Symes, Anomalous reflections near a caustic, *Wave Motion* **25** (1997), 1–14.
- [30] ———, Global solutions of a linearized inverse problem for the acoustic wave equation, *Comm. P.D.E.* **22** (1997), 919–952.
- [31] Rakesh, A linearized inverse problem for the wave equation, *Comm. P.D.E.* **13** (1988), 573–601.
- [32] A. Weinstein, On Maslov’s quantization condition, in: Chazarain, J., ed. *Fourier Integral Operators and Partial Differential Equations*, Springer-Verlag, New York, 1975.
- [33] H. Whitney, The general type of singularity of a set of $2n - 1$ smooth functions of n variables, *Duke Math. Jour.* **10** (1943), 161–172.

SCHOOL OF MATHEMATICAL SCIENCES
 ROCHESTER INSTITUTE OF TECHNOLOGY
 ROCHESTER, NY 14623
 rxfsma@rit.edu

DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF ROCHESTER
 ROCHESTER, NY 14627
 allan@math.rochester.edu